

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 October 2002 (24.10.2002)

PCT

(10) International Publication Number
WO 02/083241 A1

(51) International Patent Classification⁷: **A61P 15/00**,
5/24, A61K 31/56, 31/4196

(21) International Application Number: PCT/CA02/00527

(22) International Filing Date: 17 April 2002 (17.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/284,282 17 April 2001 (17.04.2001) US

(71) Applicant (for all designated States except US): **ARES TRADING S.A.** [CH/CH]; Chateau de Vaumarcus, CH-2028 Vaumarcus (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CASPER, Robert, F.** [CA/CA]; 9 Meredith Crescent, Toronto, Ontario M4W 3B7 (CA). **MITWALLY, Mohamed, F., M.** [EG/CA]; 387 Melrose Avenue, North York, Ontario M5M 1Z6 (CA).

(74) Agents: **VAN ZANT, Joan, M.** et al.; Ogilvy Renault, 1981 McGill College Avenue, Suite 1600, Montreal, Ontario H3A 2Y3 (CA).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/083241 A1

(54) Title: SINGLE DOSE AROMATASE INHIBITOR FOR TREATING INFERTILITY

(57) **Abstract:** The use of an aromatase inhibitor (AI) in the preparation of a medicament for use in inducing ovulation in a female suffering from anovulatory infertility which comprises administering to said female a single dose of the at least one aromatase inhibitor. The use of an aromatase inhibitor (AI) for the preparation of a medicament comprising a single dose of at least one aromatase inhibitor (AI) for administration to an ovulating female suffering from unexplained infertility or other types of ovulatory infertility early in one or more menstrual cycles. Such a medicament may also be administered to an anovulatory female in combination with a plurality of daily doses of follicle stimulating hormone (FSH) whereby the dosage levels of FSH may be reduced. The use of AI in the preparation of a medicament for administration to a female who is a poor responder to FSH, which comprises a combination of a single dose of AI with a plurality of daily doses of FSH is also disclosed. The related pharmaceutical preparations and packages are also described.

TITLE

Single Dose Aromatase Inhibitor for Treating Infertility

FIELD OF THE INVENTION

This invention relates to the use of a single dose of aromatase inhibitors (AIs) for inducing and augmenting ovulation in females with anovulatory infertility or unexplained infertility. Also described is the use of AIs together with follicle stimulating hormone (FSH) for improving response to controlled ovarian hyperstimulation. Also disclosed are pharmaceutical formulations, which comprise the aromatase inhibitors and methods for administering them to females.

10 BACKGROUND OF THE INVENTION

In women with WHO type II anovulatory infertility such as polycystic ovary syndrome (PCOS), the treatment of first choice for the induction of ovulation is an anti-estrogen. The most commonly used drug is clomiphene citrate (CC). However, 20 to 25 percent of women do not ovulate with CC. In addition, clinical data reveal a discrepancy between ovulation and conception rates during CC treatment¹, and a higher than expected incidence of miscarriage in conception cycles². These observations have been attributed to the anti-estrogenic mechanism of action of CC resulting in long-lasting estrogen receptor depletion. Thus, CC may have a negative effect on the quality and quantity of cervical mucus³, on endometrial development⁴, and on other as yet undetermined fertility factors since CC is accumulated in the body as a result of its long half-life⁵.

In CC failures, gonadotropin preparations such as human menopausal gonadotropin (hMG) or pure follicle stimulating hormone (FSH) have been used as a second-line treatment for ovulation induction. In women with polycystic ovary syndrome, because of the high sensitivity of the ovaries to gonadotropin stimulation, treatment with hMG or pure FSH is difficult to control and characteristically induces multiple follicles. The result is a high frequency of multiple pregnancies and increased risk of ovarian hyperstimulation syndrome (OHSS)⁶. Therefore, a simple oral treatment that could be used without risk of hyperstimulation and with minimal monitoring is the preferred first line of therapy.

Additionally, although it has been established that pregnancy rates for women who take CC are less than expected based on ovulation rates, CC therapy is widely administered to women with unexplained infertility, often without ultrasound monitoring, in order to induce the development of multiple follicles⁷. The use of CC in these women may be unsuccessful because of antiestrogenic effects on endometrial development. A recent study has prospectively applied morphometric analysis of the endometrium, which is a quantitative and objective technique to study the effect of CC on the endometrium in a group of normal women. In this study, CC was found to have a deleterious effect on the endometrium, demonstrated by a reduction in glandular density and an increase in the number of vacuolated cells⁸. In some exceptional cases, normal ovulatory women may receive 6 to 12 cycles of CC before it is finally determined that the anti-estrogenic effects of CC on the endometrium are actually causing an anti-conception action. For these reasons, a simple,

inexpensive and safe alternative to CC for use in normally ovulatory women, in whom frequent cycle monitoring is difficult, is also required.

The induction of ovulation constitutes a vital part of infertility management.

Unfortunately, most current therapeutic approaches for induction of ovulation have been empiric⁹. For over 40 years, clomiphene citrate (CC) has been the most commonly used treatment for the induction and augmentation of ovulation, accounting for about two thirds of the fertility drugs prescribed in the United States¹⁰. However, the mechanism(s) and site(s) of CC action have only been partially clarified despite extensive clinical research¹¹.

Mechanism of CC action

It is believed that the 2 isomers of CC exert either an anti-estrogenic effect (zu-clomiphene) or a weak estrogen agonist effect (en-clomiphene) at estrogen binding sites in the pituitary and hypothalamus, thus releasing the hypothalamic/pituitary axis from the inhibitory effect of the major circulating estrogen, estradiol (E₂)¹². In women with PCOS, CC-induced ovulation was accompanied by increased secretion of LH and FSH with enhanced estrogen secretion. Increased LH pulse amplitude after CC, together with decreased pituitary sensitivity to a GnRH bolus, suggested that CC acted predominantly on the hypothalamus to cause release of larger pulses of GnRH into the pituitary-portal system¹³. Similar findings have been reported in normal ovulatory women¹⁴. Various mechanisms of CC action have also been suggested at the level of the pituitary and/or the ovary. In particular, the ovarian actions of CC have not been widely appreciated¹⁵. However, the overall mechanism of CC action may be the sum of its effects on the hypothalamus, pituitary and ovary as discussed by Adashi¹⁶.

Approaches to improve pregnancy outcome with CC

In order to improve the outcome of CC treatment, various approaches have been suggested to overcome the antiestrogenic effect of CC. One approach has been to administer estrogen concomitantly during CC treatment to attain high estrogen levels to overcome the antiestrogenic effect of CC. Some investigators have reported success with this approach¹⁷ while others have reported no benefit¹⁸ or even a deleterious effect of estrogen administration¹⁹. Another approach to reduce adverse effects has been to administer CC earlier during the menstrual cycle rather than starting on day 5²⁰, in the hopes of allowing the anti-estrogenic effect to wear off to some extent. A third approach has been to combine another selective estrogen receptor modulator like tamoxifen, which has more estrogen agonistic effect on the endometrium with CC²¹. However, none of the above mentioned strategies have proved to be completely successful in avoiding the peripheral antiestrogenic effects of CC. In addition to a discrepancy between ovulation and pregnancy rates with CC treatment, 20% to 25% of anovulatory women are resistant to CC and fail to ovulate at doses up to 150 mg daily. In CC failures, gonadotropins have been used as a second-line treatment for ovulation induction. However, they are associated with higher risk of multiple pregnancy, and severe ovarian hyperstimulation syndrome, especially in women with PCOS. Therefore, a simple oral alternative to CC that could be

used without high risk and which requires minimal monitoring would be the preferred first line of therapy for ovulation induction.

Aromatase Inhibitor

- 5 A group of highly selective AIs has been approved for use in postmenopausal women with breast cancer to suppress estrogen production. These AIs have a relatively short half-life (approximately 48 hours) compared to CC, and therefore would be eliminated from the body rapidly²². In addition, since no estrogen receptor down-regulation occurs, no adverse effects on estrogen target tissues, as observed in CC treated cycles, would be expected.

Physiology of Aromatase Enzyme

- 10 Aromatase is a cytochrome P-450 hemoprotein-containing enzyme complex that catalyzes the rate-limiting step in the production of estrogens, i.e. the conversion of androstenedione and testosterone, via three hydroxylation steps, to estrone and estradiol²³. Aromatase activity is present in many tissues, such as the ovaries, adipose tissue, muscle, liver, breast tissue, and in malignant breast tumors. The main sources of circulating
15 estrogens are the ovaries in premenopausal women and adipose tissue in postmenopausal women²⁴. Although aromatase has features in common with other steroidogenic P-450 enzymes, the heme-binding region has only 17.9±23.5% amino acids identical to those of other steroidogenic P-450 enzymes. This observation suggests that P-450arom belongs to a separate gene family which has been designated CYP19²⁵.
20 Aromatase catalyzes the conversion of androgens to estrone (E₁), which is further converted to the potent estrogen estradiol (E₂) by the enzyme 17β-HSD type 1 in the granulosa cell.

Development of aromatase Inhibitors

- Aromatase is a good target for selective inhibition because estrogen production is a
25 terminal step in the biosynthetic sequence. There are two types of aromatase inhibitors; steroidal (type I inhibitors) and non-steroidal inhibitors (type II inhibitors). Type I steroidal aromatase inhibitors are all derivatives of androstenedione that act as a false substrate and bind irreversibly to the androgen-binding site with continuing treatment. For this reason, they are also called suicide inhibitors. 4-hydroxyandrostenedione, the first selective
30 steroidal aromatase inhibitor to be used clinically, has proved to be effective in tamoxifen-resistant breast cancer patients and is available in many countries worldwide²⁶. Type II non-steroidal aromatase inhibitors exert their function through binding to the heme moiety of the cytochrome P450 enzyme. The first of these inhibitors to be used clinically was aminoglutethimide, which induces a medical adrenalectomy by inhibiting many other
35 enzymes involved in steroid biosynthesis²⁷. Although aminoglutethimide is an effective hormonal agent in postmenopausal breast cancer, its use is complicated by the need for concurrent corticosteroid replacement, in addition to side effects like lethargy, rashes, nausea and fever that results in 8–15% of patients stopping treatment. The lack of specificity and unfavorable toxicity profile of aminoglutethimide have led to the search for

more specific aromatase inhibitors. In addition, the above mentioned aromatase inhibitors were not able to completely inhibit aromatase activity in premenopausal patients²⁸.

Aromatase inhibitors such as anastrozole, ZN 1033, (Arimidex[®]), letrozole, CGS 20267, (Femara[®]) and vorozole (Rivizor[®]) are selective AIs, available for clinical use in North America and other parts of the world for treatment of postmenopausal breast cancer. These triazole (antifungal) derivatives are reversible, competitive AIs, which are highly potent and selective (104, 106)²⁹. Their intrinsic potency is considerably greater than that of aminoglutethimide, and at doses of 1-5 mg/day, inhibit estrogen levels by 97% to >99%. This level of aromatase inhibition results in estradiol concentrations below detection by most sensitive immunoassays. The high affinity of AIs for aromatase is thought to reside in the N-4 nitrogen of the triazole ring that coordinates with the heme iron atom of the aromatase enzyme complex. AIs are completely absorbed after oral administration with mean terminal $t_{1/2}$ of approximately 50 hr (range, 30-60 hr). They are cleared from the systemic circulation mainly by the liver. Another AI available commercially is exemestane (Aromasin[™]).

In animal studies, letrozole resulted in increased FSH and LH when given to mature female rats and about a 30% increase in ovarian weight³⁰. In the bonnet monkey, treatment with aromatase inhibitors to induce estradiol deficiency led to development of multiple normal Graafian follicles in vivo, and normal response of granulosa and theca cells to gonadotropins in vitro³¹. In vivo data describe a continuum of inhibition of aromatase, with aminoglutethimide (90%), vorozole (93%), anastrozole (97%), and letrozole (98.5%) displaying increasing potency and specificity³². Letrozole has an IC_{50} of 11.5 nM in vitro and ED_{50} of 1-3 μ g/kg in vivo when given orally. The disposition of orally administered letrozole in healthy postmenopausal women is characterized by steady-state plasma concentrations in 4 to 8 hours, and a half-life of approximately 45 hours. The absolute systemic bioavailability of letrozole after oral administration was 100% compared with the same dose given intravenously³³. Doses up to 30 mg have been well tolerated³⁴. The lethal dose in mice and rats is approximately 2000 mg/kg. There is no experience in humans with an overdose of letrozole³⁵.

The success of aromatase inhibition by letrozole in inducing ovulation in women with PCOS has been reported³⁶. In a series of 10 patients with PCOS who either failed to ovulate (n=4) or ovulated with an endometrial thickness \leq 5 mm (n=6) in response to CC administration, ovulation occurred in 7 of the 10 letrozole treated cycles (70%), with clinical pregnancy in 2 patients and chemical pregnancy in one patient. The mean number of mature follicles was 2.6, ranging from 1 to 4 follicles in the 7 ovulatory cycles. The mean level of estradiol on the day of hCG administration was 1076 pmol/L with mean estradiol per follicle of 378 pmol/L. This estradiol level allowed the growth of the endometrium to an adequate thickness that ranged from 0.7 cm to 0.9cm on the day of hCG administration, showing the absence of antiestrogenic effects as seen with CC.

In a second study, comparable success of letrozole in inducing ovulation in 12 women with PCOS women, in addition to success in augmenting ovulation in a group of 10 ovulatory women. Patients in both groups tried CC in prior treatment cycles with an inadequate response. With letrozole treatment, ovulation occurred in 9 of 12 cycles (75%) and pregnancy was achieved in 3 patients (25%) in the PCOS group. In the ovulatory group, letrozole resulted in a mean number of 2.3 mature follicles and a mean endometrial thickness of 0.8 cm. Pregnancy was achieved in one patient (10%)³⁷.

We have also studied the use of letrozole in conjunction with FSH for controlled ovarian super ovulation in both ovulatory women with unexplained infertility and anovulatory women with PCOS³⁸. The use of letrozole was associated with a significantly lower FSH dose required for achievement of adequate ovarian super ovulation. The pregnancy rate and endometrial thickness with letrozole and FSH treatment was similar to FSH alone. We have also shown an improvement in ovarian response to FSH stimulation with the use of letrozole in low responders during ovarian stimulation³⁹.

In U.S. Patent No. 5,583,128 granted Dec. 10, 1996 to Bhatnagar, there is described the use of aromatase inhibitors for contraception in female primates of reproductive age without substantially affecting the menstrual cycle of the female primate. The contraceptive action of the aromatase inhibitors is reversible, that is to say once their use has been discontinued pregnancy can occur in the treated primates as early as the next cycle.

In U.S. Patent No. 5,491,136 granted Feb. 13, 1996 to Peet et al, the use of aromatase inhibitors in the treatment of breast cancer is described.

The disclosures of all references referred to herein are incorporated herein by reference.

SUMMARY OF THE INVENTION

The present invention mimics the action of CC, without depletion of estrogen receptors, by administration of an aromatase inhibitor (AI) in the early part of the menstrual cycle. This is believed to result in release of the hypothalamic/pituitary axis from estrogenic negative feedback, thereby increasing gonadotropin secretion and resulting in stimulation of ovarian follicles. In addition, a peripheral mechanism of action at the level of the ovary may also be present, secondary to prevention of androgen conversion to estrogen and an increase in intra-ovarian androgen concentration. In the primate ovary, androgen has been shown to increase granulosa cell FSH receptors^{40, 41} thereby increasing ovarian responsiveness to FSH.

The present invention provides an improvement over the known multiple daily dose administration of aromatase inhibitor for treating infertility in females. The single dose provides a high initial dose of AI with a better or equal disappearance from the body as compared with the multiple dose treatment during the critical time for follicle recruitment and stimulation. Usually this encompasses a period of about 7 days at the beginning of the menstrual cycle. The single dose is more convenient and ensures better patient compliance. The amount of aromatase inhibitor that remains in the body in the single dose

form as compared with multiple dose form has been found to be about equivalent.

Thus the present invention provides a method of inducing ovulation in a female suffering from anovulatory infertility which comprises administering to such a female a single dose of at least one aromatase inhibitor (AI).

- 5 In another form the present invention provides a method for augmenting ovulation in an ovulating female suffering from unexplained infertility or another type of ovulatory infertility, which comprises administering to such a female, a single dose of at least one aromatase inhibitor (AI), early in one or more menstrual cycles.

- 10 In another aspect the invention provides a method of substantially reducing dosage levels of follicle stimulating hormone (FSH) for administration to a female undergoing infertility treatment which comprises administering a combination of a single dose of at least one aromatase inhibitor (AI) with a plurality of daily doses of follicle stimulating hormone (FSH).

- 15 The invention also provides a method of increasing response to follicle stimulating hormone in a female who is a poor responder to follicle stimulation, which comprises administering a combination of a single dose of at least one aromatase inhibitor (AI) with a plurality of daily doses of follicle stimulating hormone (FSH).

- 20 In another aspect the invention provides a pharmaceutical preparation for treating infertility in a female comprising a single dose of an aromatase inhibitor for inducing or augmenting ovulation together with a pharmaceutically acceptable carrier

In another aspect the invention provides a two component pharmaceutical preparation for treating infertility in a female comprising a single dose of an aromatase inhibitor together with a pharmaceutically acceptable carrier in combination with a plurality of daily doses of follicle stimulating hormone together with a pharmaceutically acceptable carrier.

- 25 In another aspect the invention provides for the use of a single dose of an aromatase inhibitor for treating infertility in a female, each dose comprising an effective amount of an aromatase inhibitor for inducing or augmenting ovulation.

- 30 In another aspect the invention provides for the use of a single dose of an aromatase inhibitor in combination with a plurality of daily doses of follicle stimulating hormone for treating infertility in a female wherein the amount of follicle stimulating hormone is substantially reduced as compared with the use of follicle stimulating hormone on its own.

- 35 In another aspect the invention provides for the use a single dose of an aromatase inhibitor in combination with a plurality of daily doses of follicle stimulating hormone for treating a female who is a poor responder to follicle stimulation to increase follicle production.

In another aspect the invention provides for the use of a single dose of an aromatase inhibitor in the preparation of a medicament for the treatment of infertility in a female.

- 40 In another aspect the invention provides for the use of a single dose of an aromatase inhibitor in combination with a plurality of daily doses of follicle stimulating hormone in the preparation of a medicament for the treatment of infertility in a female.

DETAILED DESCRIPTION OF THE INVENTION

The daily doses required for the present invention depend entirely on the type of aromatase inhibitor that is used and the patient. Some inhibitors are more active than others and hence lower amounts of the former inhibitors may be used.

- 5 The aromatase inhibitor is selected from aromatase inhibitors having a half life of about 8 hours to about 4 days, more preferably from aromatase inhibitors having a half life of about 2 days. Most beneficial are those aromatase inhibitors selected from non-steroidal and reversible aromatase inhibitors. More detail on the types of aromatase inhibitors that may be used in the methods, uses and preparations of the present invention appears
10 subsequently herein.

Aromatase Inhibitor

- The aromatase inhibitors that have been found to be most useful of the commercially available forms are those in oral form. This form offers clear advantages over others, including convenience and patient compliance. Preferred aromatase inhibitors of those
15 that are commercially available include anastrozole, letrozole, vorozole and exemestane. The AI exemestane is a steroidal inhibitor.

By "aromatase inhibitors" there are to be understood substances that inhibit the enzyme aromatase (estrogen synthetase), which is responsible for converting androgens to estrogens.

- 20 Aromatase inhibitors may have a non-steroidal or a steroidal chemical structure. According to the present invention, both non-steroidal aromatase inhibitors and steroidal aromatase inhibitors can be used.

- By aromatase inhibitors there are to be understood especially those substances that in a determination of the in vitro inhibition of aromatase activity exhibit IC_{50} values of 10^{-5} M or
25 lower, especially 10^{-6} M or lower, preferably 10^{-7} M or lower and most especially 10^{-8} M or lower.


- The in vitro inhibition of aromatase activity can be demonstrated, for example, using the methods described in J. Biol. Chem. 249, 5364 (1974) or in J. Enzyme Inhib. 4, 169 (1990). In addition, IC_{50} values for aromatase inhibition can be obtained, for example, in
30 vitro by a direct product isolation method relating to inhibition of the conversion of 4- 14 C-androstenedione to 4- 14 C-oestrone in human placental microsomes.

- By aromatase inhibitors there are to be understood most especially substances for which the minimum effective dose in the case of in vivo aromatase inhibition is 10 mg/kg or less, especially 1 mg/kg or less, preferably 0.1 mg/kg or less and most especially 0.01
35 mg/kg or less.

- In vivo aromatase inhibition can be determined, for example, by the following method [see J. Enzyme Inhib. 4, 179 (1990)]: androstenedione (30 mg/kg subcutaneously) is administered on its own or together with a compound of the invention (orally or subcutaneously) to sexually immature female rats for a period of 4 days. After the fourth
40 administration, the rats are sacrificed and the uteri are isolated and weighed. The

aromatase inhibition is determined by the extent to which the hypertrophy of the uterus induced by the administration of androstenedione alone is suppressed or reduced by the simultaneous administration of the compound according to the invention.


- The following groups of compounds are listed as examples of aromatase inhibitors. Each individual group forms a group of aromatase inhibitors that can be used successfully in accordance with the present invention:
- (a) The compounds of formulae I and I* as defined in EP-A-165 904. These are especially the compounds of formula I



10

- wherein R_1 is hydrogen, lower alkyl; lower alkyl substituted by hydroxy, lower alkoxy, lower alkanoyloxy, lower alkanoyl, amino, lower alkylamino, di-lower alkylamino, halogen, sulfo, carboxy, lower alkoxy carbonyl, carbamoyl or by cyano; nitro, halogen, hydroxy, lower alkoxy, lower alkanoyloxy, phenylsulfonyloxy, lower alkylsulfonyloxy, mercapto, lower alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, lower alkanoylthio, amino, lower alkylamino, di-lower alkylamino, lower alkyleneamino, N-morpholino, N-thiomorpholino, N-piperazino that is unsubstituted or lower alkyl-substituted in the 4-position, tri-lower alkylammonio, sulfo, lower alkoxy sulfonyl, sulfamoyl, lower alkylsulfamoyl, di-lower alkylsulfamoyl, formyl; iminomethyl that is unsubstituted or substituted at the nitrogen atom by hydroxy, lower alkoxy, lower alkanoyloxy, lower alkyl, phenyl or by amino; C_2-C_7 alkanoyl, benzoyl, carboxy, lower alkoxy carbonyl, carbamoyl, lower alkylcarbamoyl, di-lower alkylcarbamoyl, cyano, 5-tetrazolyl, unsubstituted or lower alkyl-substituted 4,5-dihydro-2-oxazolyl or hydroxycarbamoyl; and R_2 is hydrogen, lower alkyl, phenyl-lower alkyl, carboxy-lower alkyl, lower alkoxy carbonyl-lower alkyl, halogen, hydroxy, lower alkoxy, lower alkanoyloxy, mercapto, lower alkylthio, phenyl-lower alkylthio, phenylthio, lower alkanoylthio, carboxy, lower alkoxy carbonyl or lower alkanoyl; the 7,8-dihydro derivatives thereof; and the compounds of formula I*

25




wherein n is 0, 1, 2, 3 or 4; and R_1 and R_2 are as defined above for formula I; it being possible for the phenyl ring in the radicals phenylsulfonyloxy, phenyliminomethyl, benzoyl, phenyl-lower alkyl, phenyl-lower alkylthio and phenylthio to be unsubstituted or substituted by lower alkyl, lower alkoxy or by halogen; it being possible in a compound of formula I* for the two substituents $C_6H_4-R_1$ and R_2 to be linked to each of the saturated carbon atoms of the saturated ring, either both to the same carbon atom or both to different carbon atoms, and pharmaceutically acceptable salts thereof.

Individual compounds that may be given special mention here are:


- (1) 5-(p-cyanophenyl)imidazo[1,5-a]pyridine,
- (2) 5-(p-ethoxycarbonylphenyl)imidazo[1,5-a]pyridine,
- (3) 5-(p-carboxyphenyl)imidazo[1,5-a]pyridine,
- (4) 5-(p-tert-butylaminocarbonylphenyl)imidazo[1,5-a]pyridine,
- (5) 5-(p-ethoxycarbonylphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (6) 5-(p-carboxyphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (7) 5-(p-carbamoylphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (8) 5-(p-tolyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (9) 5-(p-hydroxymethylphenyl)imidazo[1,5-a]pyridine,
- (10) 5-(p-cyanophenyl)-7,8-dihydroimidazo[1,5-a]pyridine,
- (11) 5-(p-bromophenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (12) 5-(p-hydroxymethylphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (13) 5-(p-formylphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (14) 5-(p-cyanophenyl)-5-methylthio-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (15) 5-(p-cyanophenyl)-5-ethoxycarbonyl-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (16) 5-(p-aminophenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (17) 5-(p-formylphenyl)imidazo[1,5-a]pyridine,
- (18) 5-(p-carbamoylphenyl)imidazo[1,5-a]pyridine,
- (19) 5H-5-(4-tert-butylaminocarbonylphenyl)-6,7-dihydropyrrolo[1,2-c]imidazole,
- (20) 5H-5-(4-cyanophenyl)-6,7-dihydropyrrolo[1,2-c]imidazole,
- (21) 5H-5-(4-cyanophenyl)-6,7,8,9-tetrahydroimidazo[1,5-a]azepine,
- (22) 5-(4-cyanophenyl)-6-ethoxycarbonylmethyl-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
- (23) 5-(4-cyanophenyl)-6-carboxymethyl-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine

- (24) 5-benzyl-5-(4-cyanophenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
 (25) 7-(p-cyanophenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
 (26) 7-(p-carbamoylphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
 (27) 5-(p-cyanophenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine (=Fadrozol).

- 5 (b) The compounds of formula I as defined in EP-A 236 940. These are especially the compounds of formula I



- 10 wherein R and R₀, independently of one another, are each hydrogen or lower alkyl, or R and R₀ at adjacent carbon atoms, together with the benzene ring to which they are bonded, form a naphthalene or tetrahydronaphthalene ring; wherein R₁ is hydrogen, lower alkyl, aryl, aryl-lower alkyl or lower alkenyl; R₂ is hydrogen, lower alkyl, aryl, aryl-lower alkyl, (lower alkyl, aryl or aryl-lower alkyl)-thio or lower alkenyl, or wherein R₁ and R₂
- 15 together are lower alkylidene or C₄-C₆ alkylene; wherein W is 1-imidazolyl, 1-(1,2,4 or 1,3,4)-triazolyl, 3-pyridyl or one of the mentioned heterocyclic radicals substituted by lower alkyl; and aryl within the context of the above definitions has the following meanings:
- 20 phenyl that is unsubstituted or substituted by one or two substituents from the group lower alkyl, lower alkoxy, hydroxy, lower alkanoyloxy, nitro, amino, halogen, trifluoromethyl, cyano, carboxy, lower alkoxy carbonyl, carbamoyl, N-lower alkylcarbamoyl, N,N-di-lower alkylcarbamoyl, lower alkanoyl, benzoyl, lower alkylsulfonyl, sulfamoyl, N-lower alkylsulfamoyl and N,N-di-lower alkylsulfamoyl; also thienyl, indolyl, pyridyl or furyl, or one of the four last-mentioned heterocyclic radicals monosubstituted by lower alkyl, lower alkoxy, cyano or by halogen; and pharmaceutically acceptable salts thereof.
- 25 Individual compounds from that group that may be given special mention are:
- (1) 4-[alpha-(4-cyanophenyl)-1-imidazolylmethyl]-benzonitrile,
 (2) 4-[alpha-(3-pyridyl)-1-imidazolylmethyl]-benzonitrile,
 (3) 4-[alpha-(4-cyanobenzyl)-1-imidazolylmethyl]-benzonitrile,
 (4) 1-(4-cyanophenyl)-1-(1-imidazolyl)-ethylene,
 30 (5) 4-[alpha-(4-cyanophenyl)-1-(1,2,4-triazolyl)methyl]-benzonitrile,
 (6) 4-[alpha-(4-cyanophenyl)-3-pyridylmethyl]-benzonitrile.
- (c) The compounds of formula I as defined in EP-A-408 509. These are especially the compounds of formula I




- wherein Tetr is 1- or 2-tetrazolyl that is unsubstituted or substituted in the 5-position by lower alkyl, phenyl-lower alkyl or by lower alkanoyl; R and R₂, independently of one another, are each hydrogen; lower alkyl that is unsubstituted or substituted by hydroxy, lower alkoxy, halogen, carboxy, lower alkoxy-carbonyl, (amino, lower alkylamino or di-lower alkylamino)-carbonyl or by cyano; lower alkenyl, aryl, heteroaryl, aryl-lower alkyl, C₃-C₆ cycloalkyl, C₃-C₆ cycloalkyl-lower alkyl, lower alkylthio, arylthio or aryl-lower alkylthio; or R₁ and R₂ together are straight-chained C₄-C₆ alkylene that is unsubstituted or substituted by lower alkyl, or are a group -(CH₂)_m-1,2-phenylene-(CH₂)_n-- wherein m and n, independently of one another, are each 1 or 2 and 1,2-phenylene is unsubstituted or substituted in the same way as phenyl in the definition of aryl below, or are lower alkylidene that is unsubstituted or mono- or di-substituted by aryl; and R and R₀, independently of one another, are each hydrogen or lower alkyl; or R and R₀ together, located at adjacent carbon atoms of the benzene ring, are a benzo group that is unsubstituted or substituted in the same way as phenyl in the definition of aryl below; aryl in the above definitions being phenyl that is unsubstituted or substituted by one or more substituents from the group consisting of lower alkyl, lower alkoxy, hydroxy, lower alkanoyloxy, nitro, amino, halogen, trifluoromethyl, carboxy, lower alkoxy-carbonyl, (amino, lower alkylamino or di-lower alkylamino)-carbonyl, cyano, lower alkanoyl, benzoyl, lower alkylsulfonyl and (amino, lower alkylamino or di-lower alkylamino)-sulfonyl; heteroaryl in the above definitions being an aromatic heterocyclic radical from the group consisting of pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, furanyl, thienyl, isoxazolyl, oxazolyl, oxadiazolyl, isothiazolyl, thiazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidyl, pyrazinyl, triazinyl, indolyl, isoindolyl, benzimidazolyl, benzotriazolyl, benzofuranyl, benzothienyl, benzoxazolyl, benzothiazolyl, benzoxadiazolyl, benzothiadiazolyl, quinolyl and isoquinolyl that is unsubstituted or substituted in the same way as phenyl in the definition of aryl above; and pharmaceutically acceptable salts thereof.

Individual compounds from that group that may be given special mention are:

- (1) 4-(2-tetrazolyl)methyl-benzonitrile,
- (2) 4-[α-(4-cyanophenyl)-(2-tetrazolyl)methyl]-benzonitrile,
- (3) 1-cyano-4-(1-tetrazolyl)methyl-naphthalene,
- (4) 4-[α-(4-cyanophenyl)-(1-tetrazolyl)methyl]-benzonitrile.


(d) The compounds of formula I as defined in European Patent Application No. 91810110.6. These are especially the compounds of formula I



- 5 wherein X is halogen, cyano, carbamoyl, N-lower alkylcarbamoyl, N-cycloalkyl-lower alkylcarbamoyl, N,N-di-lower alkylcarbamoyl, N-arylcarbamoyl, hydroxy, lower alkoxy, aryl-lower alkoxy or aryloxy, wherein aryl is phenyl or naphthyl, each of which is unsubstituted or substituted by lower alkyl, hydroxy, lower alkoxy, halogen and/or by trifluoromethyl; Y is
- 10 a group $-\text{CH}_2-\text{A}$ wherein A is 1-imidazolyl, 1-(1,2,4-triazolyl), 1-(1,3,4-triazolyl), 1-(1,2,3-triazolyl), 1-(1,2,5-triazolyl), 1-tetrazolyl or 2-tetrazolyl, or Y is hydrogen, R_1 and R_2 , independently of one another, are each hydrogen, lower alkyl or a group $-\text{CH}_2-\text{A}$ as defined for Y, or R_1 and R_2 together are $-(\text{CH}_2)_n-$ wherein n is 3, 4 or 5, with the proviso that one of the radicals Y, R_1 and R_2 is a group $-\text{CH}_2-\text{A}$, with the further proviso that in a
- 15 group $-\text{CH}_2-\text{A}$ as a meaning of R_1 or R_2 , A is other than 1-imidazolyl when X is bromine, cyano or carbamoyl, and with the proviso that in a group $-\text{CH}_2-\text{A}$ as a meaning of Y, A is other than 1-imidazolyl when X is halogen or lower alkoxy, R_1 is hydrogen and R_2 is hydrogen or lower alkyl, and pharmaceutically acceptable salts thereof.


Individual compounds from that group that may be given special mention are:

- 20 (1) 7-cyano-4-[1-(1,2,4-triazolyl)methyl]-2,3-dimethylbenzofuran,
 (2) 7-cyano-4-(1-imidazolylmethyl)-2,3-dimethylbenzofuran,
 (3) 7-carbamoyl-4-(1-imidazolylmethyl)-2,3-dimethylbenzofuran,
 (4) 7-N-(cyclohexylmethyl)carbamoyl-4-(1-imidazolylmethyl)-2,3-dimethylbenzofuran.
 (e) The compounds of formula I as defined in Swiss Patent Application 1339/90-7.
- 25 These are especially the compounds of formula I



- wherein the dotted line denotes an additional bond or no additional bond, Az is imidazolyl,
- 30 triazolyl or tetrazolyl bonded via a ring nitrogen atom, each of those radicals being

- unsubstituted or substituted at carbon atoms by lower alkyl or by aryl-lower alkyl, Z is carboxy, lower alkoxy, carbamoyl, N-lower alkylcarbamoyl, N,N-di-lower alkylcarbamoyl, N-arylcarbamoyl, cyano, halogen, hydroxy, lower alkoxy, aryl-lower alkoxy, aryloxy, lower alkyl, trifluoromethyl or aryl-lower alkyl, and R_1 and R_2 , independently of one another, are each hydrogen, lower alkyl, lower alkoxy, hydroxy, halogen or trifluoromethyl; aryl being phenyl or naphthyl each of which is unsubstituted or substituted by one or two substituents from the group consisting of lower alkyl, lower alkoxy, hydroxy, halogen and trifluoromethyl; with the proviso that neither Z nor R_2 is hydroxy in the 8-position, and pharmaceutically acceptable salts thereof.
- 10 Individual compounds from that group that may be given special mention are:
- (1) 6-cyano-1-(1-imidazolyl)-3,4-dihydronaphthalene,
 - (2) 6-cyano-1-[1-(1,2,4-triazolyl)]-3,4-dihydronaphthalene,
 - (3) 6-chloro-1-(1-imidazolyl)-3,4-dihydronaphthalene,
 - (4) 6-bromo-1-(1-imidazolyl)-3,4-dihydronaphthalene.
- 15 (f) The compounds of formula I as defined in Swiss Patent Application 3014/90-0. These are especially the compounds of formula I




- 20 wherein Z is a five-membered nitrogen-containing heteroaromatic ring selected from the group 5-isothiazolyl, 5-thiazolyl, 5-isoxazolyl, 5-oxazolyl, 5-(1,2,3-thiadiazolyl), 5-(1,2,3-oxadiazolyl), 3-(1,2,5-thiadiazolyl), 3-(1,2,5-oxadiazolyl), 4-isothiazolyl, 4-isoxazolyl, 4-(1,2,3-thiadiazolyl), 4-(1,2,3-oxadiazolyl), 2-(1,3,4-thiadiazolyl), 2-(1,3,4-oxadiazolyl), 5-(1,2,4-thiadiazolyl) and 5-(1,2,4-oxadiazolyl); R and R_0 are hydrogen; or R and R_0 together are a benzo group that is unsubstituted or substituted by lower alkyl, lower alkoxy, hydroxy, halogen or by trifluoromethyl; R_1 is hydrogen, hydroxy, chlorine or fluorine; R_3 is hydrogen; R_2 is hydrogen, lower alkyl or phenyl that is unsubstituted or substituted by lower alkyl, lower alkoxy, hydroxy, halogen, trifluoromethyl or by cyano; or R_1 and R_2 together are methyldene; or R_2 and R_3 together are $-(CH_2)_3-$; or R_1 and R_2 and R_3 together are a group $=CH-(CH_2)_2-$ wherein the single bond is linked to the benzene ring; X is cyano; and X may also be halogen when R_2 and R_3 together are $-(CH_2)_3-$ or R_1 and R_2 and R_3 together are a group $=CH-(CH_2)_2-$; and pharmaceutically acceptable salts thereof.

Individual compounds from that group that may be given special mention are:

- (1) 4-[α -(4-cyanophenyl)- α -hydroxy-5-isothiazolylmethyl]-benzonitrile.
- 35 (2) 4-[α -(4-cyanophenyl)-5-isothiazolylmethyl]-benzonitrile,


- (3) 4-[α -(4-cyanophenyl)-5-thiazolylmethyl]-benzonitrile,
 (4) 1-(4-cyanophenyl)-1-(5-thiazolyl)-ethylene,
 (5) 6-cyano-1-(5-isothiazolyl)-3,4-dihydronaphthalene,
 (6) 6-cyano-1-(5-thiazolyl)-3,4-dihydronaphthalene.
- 5 (g) The compounds of formula VI as defined in Swiss Patent Application 3014/90-0.
 These are especially the compounds of formula VI



- wherein Z is a five-membered nitrogen-containing heteroaromatic ring selected from the
- 10 group 5-isothiazolyl, 5-thiazolyl, 5-isoxazolyl, 5-oxazolyl, 5-(1,2,3-thiadiazolyl), 5-(1,2,3-oxadiazolyl), 3-(1,2,5-thiadiazolyl), 3-(1,2,5-oxadiazolyl), 4-isothiazolyl, 4-isoxazolyl, 4-(1,2,3-thiadiazolyl), 4-(1,2,3-oxadiazolyl), 2-(1,3,4-thiadiazolyl), 2-(1,3,4-oxadiazolyl), 5-(1,2,4-thiadiazolyl) and 5-(1,2,4-oxadiazolyl); R and R₀ are each hydrogen; or R and R₀ together are a benzo group that is unsubstituted or substituted by lower alkyl, lower alkoxy,
- 15 hydroxy, halogen or by trifluoromethyl; R₁ is hydrogen, hydroxy, chlorine or fluorine; R₃ is hydrogen; R₂ is hydrogen, lower alkyl or phenyl that is unsubstituted or substituted by lower alkyl, lower alkoxy, hydroxy, halogen, trifluoromethyl, aryl-lower alkoxy or by aryloxy; or R₁ and R₂ together are methyldiene, and W₂ is halogen, hydroxy, lower alkoxy, aryl-lower alkoxy or aryloxy; aryl in each case being phenyl that is
- 20 unsubstituted or substituted by lower alkyl, lower alkoxy, hydroxy, halogen or by trifluoromethyl; and pharmaceutically acceptable salts thereof.

Individual compounds from that group that may be given special mention are:

- (1) bis(4,4'-bromophenyl)-(5-isothiazolyl)methanol,
 (2) bis(4,4'-bromophenyl)-(5-isothiazolyl)methane,
 25 (3) bis(4,4'-bromophenyl)-(5-thiazolyl)methanol,
 (4) bis(4,4'-bromophenyl)-(5-thiazolyl)methane,
 (h) The compounds of formula I as defined in Swiss Patent Application 3923/90-4.
 These are especially the compounds of formula I



wherein Z is imidazolyl, triazolyl, tetrazolyl, pyrrolyl, pyrazolyl, indolyl, isoindolyl, benzimidazolyl, benzopyrazolyl, benzotriazolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazinyl, quinolinyl or isoquinolinyl, all those radicals being bonded via their heterocyclic rings and all those radicals being unsubstituted or substituted by lower alkyl, hydroxy, lower alkoxy, halogen or by trifluoromethyl; R₁ and R₂, independently of one another, are each

5 hydrogen or lower alkyl; or R₁ and R₂ together are C₃-C₄ alkylene, or a benzo group that is unsubstituted or substituted as indicated below for aryl; R is hydrogen, lower alkyl, aryl or heteroaryl, and X is cyano, carbamoyl, N-lower alkylcarbamoyl, N,N-di-lower alkylcarbamoyl, N,N-lower alkylenecarbamoyl; N,N-lower alkylenecarbamoyl interrupted by

10 --O--, --S-- or --NR"--, wherein R" is hydrogen, lower alkyl or lower alkanoyl; N-cycloalkylcarbamoyl, N-(lower alkyl-substituted cycloalkyl)-carbamoyl, N-cycloalkyl-lower alkylcarbamoyl, N-(lower alkyl-substituted cycloalkyl)-lower alkylcarbamoyl, N-aryl-lower alkylcarbamoyl, N-arylcarbamoyl, N-hydroxycarbamoyl, hydroxy, lower alkoxy, aryl-lower alkoxy or aryloxy; and wherein X is also halogen when Z is imidazolyl, triazolyl, tetrazolyl, pyrrolyl, pyrazolyl, indolyl, isoindolyl, benzimidazolyl, benzopyrazolyl or benzotriazolyl;

15 wherein aryl is phenyl or naphthyl, these radicals being unsubstituted or substituted by from 1 to 4 substituents from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower alkylene (linked to two adjacent carbon atoms), C₃-C₈ cycloalkyl, phenyl-lower alkyl, phenyl; lower alkyl that is substituted in turn by hydroxy, lower alkoxy, phenyl-lower alkoxy, lower alkanoyloxy, halogen, amino, lower alkylamino, di-lower alkylamino, mercapto, lower alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, carboxy, lower alkoxycarbonyl, carbamoyl, N-lower alkylcarbamoyl, N,N-di-lower alkylcarbamoyl and/or by cyano; hydroxy; lower alkoxy, halo-lower alkoxy, phenyl-lower alkoxy, phenoxy, lower alkenyloxy, halo-lower alkenyloxy, lower alkynyloxy, lower alkylenedioxy (linked to two adjacent carbon atoms),

20 lower alkanoyloxy, phenyl-lower alkanoyloxy, phenylcarbonyloxy, mercapto, lower alkylthio, phenyl-lower alkylthio, phenylthio, lower alkylsulfinyl, phenyl-lower alkylsulfinyl, phenylsulfinyl, lower alkylsulfonyl, phenyl-lower alkylsulfonyl, phenylsulfonyl, halogen, nitro, amino, lower alkylamino, C₃-C₈ cycloalkylamino, phenyl-lower alkylamino, phenylamino, di-lower alkylamino, N-lower alkyl-N-phenylamino, N-lower alkyl-N-phenyl-lower alkylamino;

25 lower alkyleneamino or lower alkyleneamino interrupted by --O--, --S-- or --NR"-- (wherein R" is hydrogen, lower alkyl or lower alkanoyl); lower alkanoylamino, phenyl-lower alkanoylamino, phenylcarbonylamino, lower alkanoyl, phenyl-lower alkanoyl, phenylcarbonyl, carboxy, lower alkoxycarbonyl, carbamoyl, N-lower alkylcarbamoyl, N,N-di-lower alkylcarbamoyl, N,N-lower alkylenecarbamoyl; N,N-lower alkylenecarbamoyl

30 interrupted by --O--, --S-- or --NR"--, wherein R" is hydrogen, lower alkyl or lower alkanoyl; N-cycloalkylcarbamoyl, N-(lower alkyl-substituted cycloalkyl)-carbamoyl, N-cycloalkyl-lower alkylcarbamoyl, N-(lower alkyl-substituted cycloalkyl)-lower alkylcarbamoyl, N-hydroxycarbamoyl, N-phenyl-lower alkylcarbamoyl, N-phenylcarbamoyl, cyano, sulfo, lower alkoxysulfonyl, sulfamoyl, N-lower alkylsulfamoyl, N,N-di-lower alkylsulfamoyl and N-phenylsulfamoyl; the phenyl groups

40


occurring in the substituents of phenyl and naphthyl in turn being unsubstituted or substituted by lower alkyl, lower alkoxy, hydroxy, halogen and/or by trifluoromethyl; wherein heteroaryl is indolyl, isoindolyl, benzimidazolyl, benzopyrazolyl, benzotriazolyl, benzo[b]furanyl, benzo[b]thienyl, benzoxazolyl or benzothiazolyl, those radicals being
 5 unsubstituted or substituted by from 1 to 3 identical or different substituents selected from lower alkyl, hydroxy, lower alkoxy, halogen, cyano and trifluoromethyl; and pharmaceutically acceptable salts thereof.

Those compounds are especially the compounds of formula I whereto Z is 1-imidazolyl, 1-(1,2,4-triazolyl), 1-(1,3,4-triazolyl), 1-(1,2,3-triazolyl), 1-tetrazolyl, 2-
 10 tetrazolyl, 3-pyridyl, 4-pyridyl, 4-pyrimidyl, 5-pyrimidinyl or 2-pyrazinyl; R₁ and R₂, independently of one another, are each hydrogen or lower alkyl; or R₁ and R₂ together are 1,4-butylene or a benzo group; R is lower alkyl; phenyl that is unsubstituted or substituted by cyano, carbamoyl, halogen, lower alkyl, trifluoromethyl, hydroxy, lower alkoxy or by phenoxy; or benzotriazolyl or benzo[b]furanyl, the last two radicals being unsubstituted or
 15 substituted by from 1 to 3 identical or different substituents selected from lower alkyl, halogen and cyano; and X is cyano or carbamoyl; and wherein X is also halogen when Z is 1-imidazolyl, 1-(1,2,4-triazolyl), 1-(1,3,4-triazolyl), 1-(1,2,3-triazolyl), 1-tetrazolyl 2-tetrazolyl; and pharmaceutically acceptable salts thereof.

Individual compounds that may be given special mention here are:

- 20 (1) 4-[4-(4-cyanophenyl)-α-fluoro-1-(1,2,4-triazolyl)methyl]-benzonitrile,
- (2) 4-[4-(4-cyanophenyl)-α-fluoro-(2-tetrazolyl)methyl]-benzonitrile,
- (3) 4-[4-(4-cyanophenyl)-α-fluoro-(1-tetrazolyl)methyl]-benzonitrile,
- (4) 4-[4-(4-cyanophenyl)-α-fluoro-(1-imidazolyl)methyl]-benzonitrile,
- (5) 1-methyl-6-[4-(4-chlorophenyl)-α-fluoro-1-(1,2,4-triazolyl)methyl]-benzotriazole,
- 25 (6) 4-[4-(4-cyanophenyl)-α-fluoro-1-(1,2,3-triazolyl)methyl]-benzo nitrile,
- (7) 7-cyano-4-[4-(4-cyanophenyl)-α-fluoro-1-(1,2,4-triazolyl)methyl]-2,3-dimethylbenzo[b]furan,
- (8) 4-[4-(4-bromophenyl)-α-fluoro-1-(1,2,4-triazolyl)methyl]-benzo nitrile,
- (9) 4-[4-(4-cyanophenyl)-α-fluoro-(5-pyrimidyl)methyl]-benzonitrile,
- 30 (10) 4-[4-(4-bromophenyl)-α-fluoro-(5-pyrimidyl)methyl]-benzonitrile,
- (11) 4-[4-(4-cyanophenyl)-α-fluoro-(3-pyridyl)methyl]-benzonitrile,
- (12) 7-bromo-4-[4-(4-cyanophenyl)-α-fluoro-(1-imidazolyl)methyl]-2,3-dimethylbenzo[b]furan,
- (13) 7-bromo-4-[4-(4-cyanophenyl)-α-fluoro-1-(1,2,4-triazolyl)methyl]-2,3-dimethylbenzo[b]furan,
- 35 (14) 4-[4-(4-cyanophenyl)-α-fluoro-(5-pyrimidyl)methyl]-benzonitrile,
- (15) 4-[4-(4-bromophenyl)-α-fluoro-(5-pyrimidyl)methyl]-benzonitrile,
- (16) 4-[4-(4-cyanophenyl)-1-(1,2,3-triazolyl)methyl]-benzonitrile,
- (17) 2,3-dimethyl-4-[4-(4-cyanophenyl)-1-(1,2,4-triazolyl)methyl]-7-cyano-benzo[b]furan,


- (18) 4-[α -(4-cyanophenyl)-(5-pyrimidyl)methyl]-benzonitrile,
 (19) 4-[α -(4-bromophenyl)-(5-pyrimidyl)methyl]-benzonitrile,
 (20) 2,3-dimethyl-4-[α -(4-cyanophenyl)-(1-imidazolyl)methyl]-7-bromo-benzo[b]furan,
 (21) 2,3-dimethyl-4-[α -(4-cyanophenyl)-1-(1,2,4-triazolyl)methyl]-7-bromo-benzo[b]furan.
- 5 (i) The compounds of formula I as defined in EP-A-114 033. These are especially the compounds of formula I



- 10 wherein R_1 is hydrogen, R_2 is hydrogen, sulfo, C_1 - C_7 alkanoyl or C_1 - C_7 alkanesulfonyl and R_3 is hydrogen, or wherein R_1 is C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_7 alkynyl, C_3 - C_{10} cycloalkyl, C_3 - C_{10} cycloalkenyl, C_3 - C_6 cycloalkyl- C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl- C_2 - C_4 alkenyl or C_3 - C_6 cycloalkenyl- C_1 - C_4 alkyl, R_2 is hydrogen, C_1 - C_7 alkyl, sulfo, C_1 - C_7 alkanoyl or C_1 - C_7 alkanesulfonyl and R_3 is hydrogen or C_1 - C_7 alkyl, and salts of those
 15 compounds.

Individual compounds from that group that may be given special mention are:

- (1) 1-(4-aminophenyl)-3-methyl-3-azabicyclo[3.1.0]hexane-2,4-dione,
 (2) 1-(4-aminophenyl)-3-n-propyl-3-azabicyclo[3.1.0]hexane-2,4-dione,
 (3) 1-(4-aminophenyl)-3-isobutyl-3-azabicyclo[3.1.0]hexane-2,4-dione,
 20 (4) 1-(4-aminophenyl)-3-n-heptyl-3-azabicyclo[3.1.0]hexane-2,4-dione,
 (5) 1-(4-aminophenyl)-3-cyclohexylmethyl-3-azabicyclo[3.1.0]hexane-2,4-dione.
- (j) The compounds of formula I as defined in EP-A-166 692. These are especially the compounds of formula I




25

- wherein R_1 is hydrogen, alkyl having from 1 to 12 carbon atoms, alkenyl having from 2 to 12 carbon atoms, lower alkynyl, cycloalkyl or cycloalkenyl each having from 3 to 10 carbon atoms, cycloalkyl-lower alkyl having from 4 to 10 carbon atoms, cycloalkyl-lower alkenyl
 30 having from 5 to 10 carbon atoms, cycloalkenyl-lower alkyl having from 4 to 10 carbon

- atoms, or aryl having from 6 to 12 carbon atoms or aryl-lower alkyl having from 7 to 15 carbon atoms, each of which is unsubstituted or substituted by lower alkyl, hydroxy, lower alkoxy, acyloxy, amino, lower alkylamino, di-lower alkylamino, acylamino amino or by halogen, R_2 is hydrogen, lower alkyl, sulfo, lower alkanoyl or lower alkanesulfonyl, sulfonyl, R_3 is hydrogen or lower alkyl and R_4 is hydrogen, lower alkyl, phenyl or phenyl substituted by $-N(R_2)(R_3)$, and salts thereof, radicals described as "lower" containing up to and including 7 carbon atoms.

Individual compounds from that group that may be given special mention are:


- (1) 1-(4-aminophenyl)-3-n-propyl-3-azabicyclo[3.1.1]heptane-2,4-dione,
 - (2) 1-(4-aminophenyl)-3-methyl-3-azabicyclo[3.1.1]heptane-2,4-dione,
 - (3) 1-(4-aminophenyl)-3-n-decyl-3-azabicyclo[3.1.1]heptane-2,4-dione,
 - (4) 1-(4-aminophenyl)-3-cyclohexyl-3-azabicyclo[3.1.1]heptane-2,4-dione,
 - (5) 1-(4-aminophenyl)-3-cyclohexylmethyl-3-azabicyclo[3.1.1]heptane-2,4-dione.
- (k) The compounds of formula I as defined in EP-A-356 673. These are especially the compounds of formula I



- wherein W (α) is a 2-naphthyl or 1-anthryl radical, wherein each benzene ring is unsubstituted or substituted by a substituent selected from halogen, hydroxy, carboxy, cyano and nitro; or (β) is 4-pyridyl, 2-pyrimidyl or 2-pyrazinyl, each of those radicals being unsubstituted or substituted by a substituent selected from halogen, cyano, nitro, $C_1 - C_4$ alkoxy and $C_2 - C_5$ alkoxycarbonyl; and pharmaceutically acceptable salts thereof.

Individual compounds from that group that may be given special mention are:


- (1) 5-(2'-naphthyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine,
 - (2) 5-(4'-pyridyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine.
- (l) The compounds of formula I or Ia as defined in EP-A-337 929. These are especially the compounds of formula I/Ia



wherein R_1 is hydrogen, methyl, ethyl, propyl, propenyl, isopropyl, butyl, hexyl, octyl, decyl, cyclopentyl, cyclohexyl, cyclopentylmethyl, cyclohexylmethyl or benzyl, R_2 is benzyloxy, 3-bromo-, 4-bromo-, 4-chloro-, 2,3-, 2,4-, 4,5- or 4,6-dichloro-benzyloxy, and R_3 is cyano; C_2 - C_{10} alkanoyl that is unsubstituted or mono- or poly-substituted by halogen, methoxy, amino, hydroxy and/or by cyano; benzoyl that is unsubstituted or substituted by one or more substituents from the group halogen, C_1 - C_4 alkyl, methoxy, amino, hydroxy and cyano; carboxy, (methoxy, ethoxy or butoxy)-carbonyl, carbamoyl, N-isopropylcarbamoyl, N-phenylcarbamoyl, N-pyrrolidylcarbonyl, nitro or amino; and salts thereof.

Individual compounds from that group that may be given special mention are:

- (1) 4-(2,4-dichlorobenzyloxy)-3-[1-(1-imidazolyl)-butyl]-benzonitrile,
 - (2) 4-(4-bromobenzyloxy)-3-[1-(1-imidazolyl)-butyl]-phenyl pentyl ketone,
 - (3) 4-(4-bromobenzyloxy)-3-[1-(1-imidazolyl)-butyl]-benzanilide,
 - (4) 4-(4-bromobenzyloxy)-3-[1-(1-imidazolyl)-butyl]-benzoic acid,
 - (5) 3-(2,4-dichlorobenzyloxy)-4-[1-(1-imidazolyl)-butyl]-benzonitrile,
 - (6) 3-(2,4-dichlorobenzyloxy)-4-[1-(1-imidazolyl)-butyl]-benzoic acid methyl ester,
 - (7) 3-(2,4-dichlorobenzyloxy)-4-[1-(1-imidazolyl)-butyl]-benzoic acid,
 - (8) 3-(3-bromobenzyloxy)-4-[1-(1-imidazolyl)-butyl]-benzonitrile,
 - (9) 4-(3-bromobenzyloxy)-3-[1-(1-imidazolyl)-butyl]-benzonitrile,
 - (10) 3-(4-bromobenzyloxy)-4-[1-(1-imidazolyl)-butyl]-benzoic acid,
 - (11) 3-(4-bromobenzyloxy)-4-[1-(1-imidazolyl)-butyl]-benzanilide,
 - (12) 3-(4-bromobenzyloxy)-4-[1-(1-imidazolyl)-butyl]-phenyl pentyl ketone,
 - (13) 4-(4-bromobenzyloxy)-3-[1-(1-imidazolyl)-butyl]-benzonitrile,
 - (14) 3-(4-bromobenzyloxy)-4-[1-(1-imidazolyl)-butyl]-benzonitrile,
 - (15) 4-nitro-2-[1-(1-imidazolyl)-butyl]-phenyl-(2,4-dichlorobenzyl) ether,
 - (16) 4-amino-2-[1-(1-imidazolyl)-butyl]-phenyl-(2,4-dichlorobenzyl) ether,
 - (17) (2,4-dichlorobenzyl)-[2-(1-imidazolyl-methyl)-4-nitrophenyl]ether.
- (m) The compounds of formula I as defined in EP-A-337 928. These are especially the compounds of formula I



30

wherein R_1 is hydrogen, methyl, ethyl, propyl, propenyl, isopropyl, butyl, hexyl, octyl, decyl, cyclopentyl, cyclohexyl, cyclopentylmethyl, cyclohexylmethyl or benzyl, R_2 is hydrogen, halogen, cyano, methyl, hydroxymethyl, cyanomethyl, methoxymethyl, pyrrolidinylmethyl, carboxy, (methoxy, ethoxy or butoxy)-carbonyl, carbamoyl, N-isopropylcarbamoyl, N-phenylcarbamoyl, N-pyrrolidylcarbonyl; C_2 - C_{10} alkanoyl that is


35

unsubstituted or mono- or poly-substituted by halogen, methoxy, ethoxy, amino, hydroxy and/or by cyano; or benzoyl that is unsubstituted or substituted by one or more substituents from the group halogen, C₁-C₄ alkyl, methoxy, ethoxy, amino, hydroxy and cyano, R₃ is hydrogen, benzyloxy, 3-bromo-, 4-bromo-, 4-chloro-, 2,3-, 2,4-, 4,5- or 4,6-

5 dichlorobenzyloxy, and X is --CH=N--; --CH=N(--O)-- or --S--; and salts thereof.

Individual compounds from that group that may be given special mention are:

- (1) 5-[1-(1-imidazolyl)-butyl]-thiophene-2-carbonitrile,
- (2) 2-[1-(1-imidazolyl)-butyl]-thiophene-4-carbonitrile,
- (3) 2-[1-(1-imidazolyl)-butyl]-4-bromo-thiophene,
- 10 (4) 2-[1-(1-imidazolyl)-butyl]-5-bromo-thiophene,
- (5) 5-[1-(1-imidazolyl)-butyl]-2-thienyl pentyl ketone,
- (6) 5-[1-(1-imidazolyl)-butyl]-2-thienyl ethyl ketone,
- (7) 5-(4-chlorobenzyloxy)-4-[1-(1-imidazolyl)-pentyl]-pyridine-2-carbonitrile,
- (8) 3-(4-chlorobenzyloxy)-4-[1-(1-imidazolyl)-pentyl]-pyridine-2-carbonitrile,
- 15 (9) 3-(4-chlorobenzyloxy)-4-[1-(1-imidazolyl)-pentyl]-pyridine-N-oxide,
- (10) 3-(4-chlorobenzyloxy)-4-[1-(1-imidazolyl)-pentyl]-pyridine.
- (n) The compounds of formula I as defined in EP-A-340 153. These are especially the compounds of formula I



20

wherein R₁ is hydrogen, methyl, ethyl, propyl, propenyl, isopropyl, butyl, hexyl, octyl, decyl, cyclopentyl, cyclohexyl, cyclopentylmethyl, cyclohexylmethyl or benzyl, and R₂ is a radical from the group methyl, ethyl, propyl, benzyl, phenyl and ethenyl that is substituted by


25 hydroxy, cyano, methoxy, butoxy, phenoxy, amino, pyrrolidinyl, carboxy, lower alkoxy carbonyl or by carbamoyl; or R₂ is formyl or derivatised formyl that can be obtained by reaction of the formyl group with an amine or amine derivative from the group hydroxylamine, O-methylhydroxylamine, O-ethylhydroxylamine, O-allylhydroxylamine, O-benzylhydroxylamine, O-4-nitrobenzyloxyhydroxylamine, O-2,3,4,5,6-

30 pentafluorobenzyloxyhydroxylamine, semicarbazide, thiosemicarbazide, ethylamine and aniline; acetyl, propionyl, butyryl, valeryl, caproyl; benzoyl that is unsubstituted or substituted by one or more substituents from the group halogen, C₁-C₄-alkyl, methoxy, amino, hydroxy and cyano; carboxy, (methoxy, ethoxy or butoxy)carbonyl, carbamoyl, N-isopropylcarbamoyl, N-phenylcarbamoyl or N-pyrrolidylcarbonyl; and salts thereof.


35 Individual compounds from that group that may be given special mention are:

- (1) 4-(1-(1-imidazolyl)-butyl)-benzoic acid methyl ester,

- (2) 4-(1-(1-imidazolyl)-butyl)-benzoic acid butyl ester,
 (3) 4-(1-(1-imidazolyl)-butyl)-phenyl-acetonitrile,
 (4) 4-(1-(1-imidazolyl)-butyl)-benzaldehyde,
 (5) 4-(1-(1-imidazolyl)-butyl)-benzyl alcohol,
 5 (6) {4-[1-(1-imidazolyl)-butyl]-phenyl }-2-propyl ketone,
 (7) 4-[1-(1-imidazolyl)-butyl]-phenyl propyl ketone,
 (8) 4-[1-(1-imidazolyl)-butyl]-phenyl butyl ketone,
 (9) 4-[1-(1-imidazolyl)-butyl]-phenyl pentyl ketone,
 (10) 4-[1-(1-imidazolyl)-butyl]-phenyl hexyl ketone.
 10 (o) The compounds of formula I as defined in DE-A-4 014 006. These are especially the compounds of formula I



- 15 wherein A is an N-atom or a CH radical and W is a radical of the formula




- wherein X is an oxygen or a sulfur atom or a $-\text{CH}=\text{CH}-$ group and Y is a methylene group, an oxygen or a sulfur atom and Z is a $-(\text{CH}_2)_n-$ group wherein $n=1, 2$ or 3 and either
- 20 a) R_3 in W is a hydrogen atom and R_1 and R_2 , independently of one another, are each a hydrogen atom, a C_1 -- to C_{10} alkyl group or a C_3 -- to C_7 cycloalkyl group, or
- b) R_2 is as defined under a) and R_1 together with R_3 forms a $-(\text{CH}_2)_m-$ group wherein $m=2, 3$, or 4 , and their pharmaceutically acceptable addition salts with acids.

Individual compounds from that group that may be given special mention are:

- 25 (1) 5-[1-(1-imidazolyl)-butyl]-1-indanone,
 (2) 7-[1-(1-imidazolyl)-butyl]-1-indanone,
 (3) 6-[1-(1-imidazolyl)-butyl]-1-indanone,

- (4) 6-(1-imidazolyl)-6,7,8,9-tetrahydro-1H-benz[e]inden-3(2H)-one,
 (5) 2-[1-(1-imidazolyl)-butyl]-4,5-dihydro-6-oxo-cyclopenta[b]-thiophene,
 (6) 6-[1-(1-imidazolyl)-butyl]-3,4-dihydro-2H-naphthalen-1-one,
 (7) 2-[1-(1-imidazolyl)-butyl]-6,7-dihydro-5H-benzo[b]thiophen-4-one,
 5 (8) 6-[1-(1-imidazolyl)-butyl]-2H-benzo[b]furan-3-one,
 (9) 5-[cyclohexyl-(1-imidazolyl)-methyl]-1-indanone,
 (10) 2-[1-(1-imidazolyl)-butyl]-4,5-dihydro-6H-benzo[b]thiophen-7-one,
 (11) 5-[1-(1-imidazolyl)-1-propyl-butyl]-1-indanone,
 (12) 2-[1-(1-imidazolyl)-butyl]-4,5-dihydro-6H-benzo[b]thiophen-7-one,
 10 (13) 2-[1-(1-imidazolyl)-butyl]-4,5-dihydro-6-oxo-cyclopenta[b]-thiophene,
 (14) 5-(1-imidazolylmethyl)-1-indanone,
 (15) 5-[1-(1,2,4-triazolyl)-methyl]-1-indanone.
 (p) The compounds of formula I as disclosed in DE-A-3 926 365. These are especially the
 compounds of formula I

15




- wherein W' is a cyclopentylidene, cyclohexylidene, cycloheptylidene or 2-adamantylidene
 radical, X is the grouping $-\text{CH}=\text{CH}-$, an oxygen or a sulfur atom, and Y and Z,
 20 independently of one another, are each a methine group (CH) or a nitrogen atom, and their
 pharmaceutically acceptable addition salts with acids.

Individual compounds from that group that may be given special mention are:

- (1) 4-[1-cyclohexylidene-1-(imidazolyl)-methyl]-benzonitrile,
 (2) 4-[1-cyclopentylidene-1-(imidazolyl)-methyl]-benzonitrile,
 25 (3) 4-[1-cycloheptylidene-1-(imidazolyl)-methyl]-benzonitrile,
 (4) 4-[2-adamantylidene-1-(imidazolyl)-methyl]-benzonitrile,
 (5) 4-[1-cyclohexylidene-1-(1,2,4-triazolyl)-methyl]-benzonitrile,
 (6) 4-[1-cyclopentylidene-1-(1,2,4-triazolyl)-methyl]-benzonitrile,
 (7) 4-[1-cycloheptylidene-1-(1,2,4-triazolyl)-methyl]-benzonitrile,
 30 (8) 4-[2-adamantylidene-1-(1,2,4-triazolyl)-methyl]-benzonitrile,
 (9) 4-[1-cyclohexylidene-1-(1,2,3-triazolyl)-methyl]-benzonitrile,
 (10) 4-[1-cyclopentylidene-1-(1,2,3-triazolyl)-methyl]-benzonitrile,
 (11) 5-[cyclohexylidene-1-imidazolylmethyl]-thiophene-2-carbonitrile.

(q) The compounds of formula I as defined in DE-A-3 740 125. These are especially the compounds of formula I



5


wherein X is CH or N, R₁ and R₂ are identical or different and are each phenyl or halophenyl, and R₃ is C₁-C₄ alkyl; C₁-C₄ alkyl substituted by CN, C₁-C₄ alkoxy, benzyloxy or by C₁-C₄ alkoxy-(mono-, di- or tri-)ethyleneoxy; C₁-C₄ alkoxy, phenyl; phenyl that is substituted by halogen or by cyano; a C₅-C₇ cycloalkyl group that is optionally condensed by benzene, or is thienyl, pyridyl or 2- or 3-indolyl; and acid addition salts thereof.

10

An individual compound from that group that may be given special mention is:

(1) 2,2-bis(4-chlorophenyl)-2-(1H-imidazol-1-yl)-1-(4-chlorobenzoyl-amino) ethane.

(r) The compounds of formula I as defined in EP-A-293 978. These are especially the compounds of formula I



15

pharmaceutically acceptable salts and stereochemically isomeric forms thereof, wherein -- A₁=A₂-A₃=A₄-- is a divalent radical selected from --CH=N--CH=CH--, --CH=N--CH=N-- and --CH=N--N=CH--, R is hydrogen or C₁-C₆ alkyl; R₁ is hydrogen, C₁-C₁₀ alkyl, C₃-C₇ cycloalkyl, Ar₁, Ar₂-C₁-C₆ alkyl, C₂-C₈ alkenyl or C₂-C₆ alkynyl; R₂ is hydrogen; C₁-C₁₀ alkyl that is unsubstituted or substituted by Ar₁; C₃-C₇ cycloalkyl, hydroxy, C₁-C₆ alkoxy, Ar₁, C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₃-C₇ cycloalkyl, bicyclo[2.2.1]heptan-2-yl, 2,3-dihydro-1H-indenyl, 1,2,3,4-tetrahydronaphthyl, hydroxy; C₂-C₆ alkenyloxy that is unsubstituted or substituted by Ar₂; C₂-C₆ alkynyloxy; pyrimidyloxy; di(Ar₂)methoxy, (1-C₁-C₄ alkyl-4-piperidinyl)oxy, C₁-C₁₀ alkoxy; or C₁-C₁₀ alkoxy that is substituted by halogen, hydroxy, C₁-C₆ alkoxy, amino, mono- or di-(C₁-C₆ alkyl)amino, trifluoromethyl, carboxy, C₁-C₆ alkoxycarbonyl, Ar.sub.I, Ar₂--O--, Ar₂--S--, C₃-C₇ cycloalkyl, 2,3-dihydro-1,4-benzodioxinyl, 1H-benzimidazolyl, C₁-C₄ alkyl-substituted 1H-benzimidazolyl, (1,1'-

20

25


- biphenyl)-4-yl or by 2,3-dihydro-2-oxo-1H-benzimidazolyl; and R_3 is hydrogen, nitro, amino, mono- or di-(C_1 - C_6 alkyl)amino, halogen, C_1 - C_6 alkyl, hydroxy or C_1 - C_6 alkoxy; wherein Ar_1 is phenyl, substituted phenyl, naphthyl, pyridyl, aminopyridyl, imidazolyl, triazolyl, thienyl, halothieryl, furanyl, C_1 - C_6 alkylfuranyl, halofuranyl or thiazolyl; wherein Ar_2 is phenyl, substituted phenyl or pyridyl; and wherein "substituted phenyl" is phenyl that is substituted by up to 3 substituents in each case selected independently of one another from the group consisting of halogen, hydroxy, hydroxymethyl, trifluoromethyl, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxy carbonyl, carboxy, formyl, hydroxyiminomethyl, cyano, amino, mono- and di-(C_1 - C_6 alkyl)amino and nitro.
- Individual compounds from that group that may be given special mention are:
- (1) 6-[(1H-imidazol-1-yl)-phenylmethyl]-1-methyl-1H-benzotriazole,
 - (2) 6-[(4-chlorophenyl)(1H-1,2,4-triazol-1-yl)methyl]-1-methyl-1H-benzotriazole.
 - (s) The compounds of formula II as defined in EP-A-250 198, especially
 - (1) 2-(4-chlorophenyl)-1,1-di(1,2,4-triazol-1-ylmethyl)ethanol,
 - (2) 2-(4-fluorophenyl)-1,1-di(1,2,4-triazol-1-ylmethyl)ethanol,
 - (3) 2-(2-fluoro-4-trifluoromethylphenyl)-1,1-di(1,2,4-triazol-1-ylmethyl)ethanol,
 - (4) 2-(2,4-dichlorophenyl)-1,1-di(1,2,4-triazol-1-ylmethyl)ethanol,
 - (5) 2-(4-chlorophenyl)-1,1-di(1,2,4-triazol-1-ylmethyl)-ethanol,
 - (6) 2-(4-fluorophenyl)-1,1-di(1,2,4-triazol-1-yl-methyl)ethanol.
 - (t) The compounds of formula I as defined in EP-A-281 283, especially
 - (1) (1R²,2R^{*})-6-fluoro-2-(4-fluorophenyl)-1,2,3,4-tetrahydro-1-(1H-1,2,4-triazol-1-yl-methyl)naphthalene,
 - (2) (1R^{*},2R^{*})-6-fluoro-2-(4-fluorophenyl)-1,2,3,4-tetrahydro-1-(1H-imidazolylmethyl)-naphthalene,
 - (3) (1R^{*},2R^{*})- and (1R^{*},2S^{*})-2-(4-fluorophenyl)-1,2,3,4-tetrahydro-1-(1H-1,2,4-triazol-1-ylmethyl)naphthalene-6-carbonitrile,
 - (4) (1R^{*},2R^{*})- and (1R^{*},2S^{*})-2-(4-fluorophenyl)-1,2,3,4-tetrahydro-1-(1H-imidazolylmethyl)naphthalene-6-carbonitrile,
 - (5) (1R^{*},2R^{*})- and (1R^{*},2S^{*})-1,2,3,4-tetrahydro-1-(1H-1,2,4-triazol-1-ylmethyl)-naphthalene-2,6-dicarbonitrile,
 - (6) (1R^{*},2R^{*})- and (1R^{*},2S^{*})-1,2,3,4-tetrahydro-1-(1H-imidazol-1-ylmethyl)naphthalene-2,6-dicarbonitrile,
 - (7) (1R^{*},2S^{*})-2-(4-fluorophenyl)-1,2,3,4-tetrahydro-1-(5-methyl-1H-imidazolyl-methyl)naphthalene-6-carbonitrile.
 - (u) The compounds of formula I as defined in EP-A-296 749, especially
 - (1) 2,2'-[5-(1H-1,2,4-triazol-1-ylmethyl)-1,3-phenylene]di(2-methylpropionitrile),
 - (2) 2,2'-[5-(imidazol-1-ylmethyl)-1,3-phenylene]di(2-methylpropionitrile),
 - (3) 2-[3-(1-hydroxy-1-methylethyl)-5-(5H-1,2,4-triazol-1-ylmethyl)phenyl]-2-methylpropionitrile,

- (4) 2,2'-[5-dideuterio(1H-1,2,4-triazol-1-yl)methyl-1,3-phenylene]di(2-trideuteriomethyl-3,3,3-trideuteriopropionitrile),
- (5) 2,2'-[5-dideuterio(1H-1,2,4-triazol-1-yl)methyl-3-phenylene]di(2methylpropionitrile).
- (v) The compounds of formula I as defined in EP-A-299 683, especially
- 5 (1) (Z)- α -(1,2,4-triazol-1-ylmethyl)stilbene-4,4'-dicarbonitrile,
- (2) (Z)-4'-chloro- α -(1,2,4-triazol-1-ylmethyl)stilbene-4-carbonitrile,
- (3) (Z)- α -(1,2,4-triazol-1-ylmethyl)-4'-(trifluoromethyl)stilbene-4-carbonitrile,
- (4) (E)-.beta.-fluoro- α -(1,2,4-triazol-1-ylmethyl)stilbene-4,4'-dicarbonitrile,
- (5) (Z)-4'-fluoro- α -(imidazol-1-ylmethyl)stilbene-4-carbonitrile,
- 10 (6) (Z)-2', 4'-dichloro- α -(imidazol-1-ylmethyl)stilbene-4-carbonitrile,
- (7) (Z)-4'-chloro- α -(imidazol-1-ylmethyl)stilbene-4-carbonitrile,
- (8) (Z)- α -(imidazol-1-ylmethyl)stilbene-4,4'dicarbonitrile,
- (9) (Z)- α -(5-methylimidazol-1-ylmethyl)stilbene-4,4'-dicarbonitrile,
- (10) (Z)-2-[2-(4-cyanophenyl)-3-(1,2,4-triazol-1-yl)propenyl]pyridine-5-carbonitrile.
- 15 (w) The compounds of formula I as defined in EP-A-299 684, especially
- (1) 2-(4-chlorobenzyl)-2-fluoro-1,3-di(1,2,4-triazol-1-yl)propane,
- (2) 2-fluoro-2-(2-fluoro-4-chlorobenzyl)-1,3-di(1,2,4-triazol-1-yl)propane,
- (3) 2-fluoro-2-(2-fluoro-4-trifluoromethylbenzyl)-1,3-di(1,2,4-triazol-1-yl)propane,
- (4) 3-(4-chlorophenyl)-1-(1,2,4-triazol-1-yl)-2-(1,2,4-triazol-1-ylmethyl)butan-2-ol,
- 20 (5) 2-(4-chloro- α -fluorobenzyl)-1,3-di(1,2,4-triazol-1-yl)propan-2-ol,
- (6) 2-(4-chlorobenzyl)-1,3-bis(1,2,4-triazol-1-yl)propane,
- (7) 4-[2-(4-chlorophenyl)-1,3-di(1,2,4-triazol-1-ylmethyl)ethoxymethyl]benzonitrile,
- (8) 1-(4-fluorobenzyl)-2-(2fluoro-4-trifluoromethylphenyl)-1,3-di(1,2,4-triazol-1-yl)-propan-2-ol,
- 25 (9) 2-(4-chlorophenyl)-1-(4-fluorophenoxy)-1,3-di(1,2,4-triazol-1-yl)propan-2-ol,
- (10) 1-(4-cyanobenzyl)-2-(2,4-difluorophenyl)-1,3di(1,2,4-triazol-1-yl)propan-2-ol,
- (11) 2-(4-chlorophenyl)-1-phenyl-1,3-di(1,2,4-triazol-1-yl)propan-2-ol.
- (x) The compounds as defined in claim 1 of EP-A-316 097, especially
- (1) 1,1-dimethyl-8-(1H-1,2,4-triazol-1-ylmethyl)-2(1H)-naphtho[2,1-b]furanone,
- 30 (2) 1,2-dihydro-1,1-dimethyl-2-oxo-8-(1H-1,2,4-triazol-1-ylmethyl)naphtho[2,1-b]-furan-7-carbonitrile,
- (3) 1,2-dihydro-1,1-dimethyl-2-oxo-8-(1H-1,2,4-triazol-1-ylmethyl)naphtho[2,1-b]-furan-7-carboxamide,
- (4) 1,2-dihydro-1,1-dimethyl-2-oxo-8-[di(1H-1,2,4-triazol-1-yl)methyl]naphtho[2,1-b]-furan-
- 35 7-carbonitrile.
- (y) The compounds of formula I as defined in EP-A-354 689, especially
- (1) 4-[2-(4-cyanophenyl)-3-(1,2,4-triazol-1-yl)propyl]benzonitrile,
- (2) 4-[1-(4-chlorobenzyl)-2-(1,2,4-triazol-1-yl)ethyl]benzonitrile,
- (3) 4-[2-(1,2,4-triazol-1-yl)-1-(4-trifluoromethyl)benzyl]ethyl]benzonitrile,
- 40 (4) 4-[2-(1,2,4-triazol-1-yl)-1-(4-[trifluoromethoxy]benzyl)ethyl]benzonitrile .

- (z) The compounds of formula (1) as defined in EP-A-354 683, especially
- (1) 6-[2-(4-cyanophenyl)-3-(1,2,4-triazol-1-yl)-propyl]nicotinonitrile,
- (2) 4-[1-(1,2,4-triazol-1-yl-methyl)-2-(5-[trifluoromethyl]pyrid-2-yl)ethyl]benzonitrile.

Examples of steroidal aromatase inhibitors that may be mentioned are:

- 5 (aa) The compounds of formula I as defined in EP-A-181 287. These are especially the compounds of formula I



- 10 wherein R is hydrogen, acetyl, heptanoyl or benzoyl. An individual compound from that group that may be given special mention is:
- (1) 4-hydroxy-4-androstene-3,17-dione.
- (ab) The compounds as defined in the claims of U.S. Pat. No. 4,322,416, especially 10-(2-propynyl)-oestr-4-ene-3,17-dione.
- 15 (ac) The compounds as defined in the claims of DE-A-3 622 841, especially 6-methyleneandrosta-1,4-diene-3,17-dione.
- (ad) The compounds as defined in the claims of GB-A-2 17 1100, especially 4-amino-androsta-1,4,6-triene-3,17-dione.

Also: (ae) androsta-1,4,6-triene-3,17-dione.

- 20 The content of the patent applications mentioned under (a) to (z) and (aa) to (ad), especially the subgroups of compounds disclosed therein and the individual compounds disclosed therein as examples, have been incorporated by reference into the disclosure of the present application.

- The general terms used hereinbefore and hereinafter to define the compounds have
- 25 the following meanings:

Organic radicals designated by the term "lower" contain up to and including 7, preferably up to and including 4, carbon atoms.

Acyl is especially lower alkanoyl.

- Aryl is, for example, phenyl or 1- or 2-naphthyl, each of which is unsubstituted or
- 30 substituted by lower alkyl, hydroxy, lower alkoxy, lower alkanoyloxy, amino, lower alkylamino, di-lower alkylamino, lower alkanoylamino or by halogen.

Pharmaceutically acceptable salts of the above-mentioned compounds are, for example, pharmaceutically acceptable acid addition salts or pharmaceutically acceptable metal or ammonium salts.

Pharmaceutically acceptable acid addition salts are especially those with suitable inorganic or organic acids, for example strong mineral acids, such as hydrochloric acid, sulfuric acid or phosphoric acid, or organic acids, especially aliphatic or aromatic carboxylic or sulfonic acids, for example formic, acetic, propionic, succinic, glycolic, lactic, 5 hydroxysuccinic, tartaric, citric, maleic, fumaric, hydroxymaleic, pyruvic, phenylacetic, benzoic, 4-aminobenzoic, anthranilic, 4-hydroxybenzoic, salicylic, 4-aminosalicylic, pamoic, gluconic, nicotinic, methanesulfonic, ethanesulfonic, halobenzenesulfonic, p-toluenesulfonic, naphthalenesulfonic, sulfanilic or cyclohexylsulfamic acid; or with other acidic organic substances, for example ascorbic acid. Pharmaceutically acceptable salts 10 may also be formed, for example, with amino acids, such as arginine or lysine.

Compounds containing acid groups, for example a free carboxy or sulfo group, can also form pharmaceutically acceptable metal or ammonium salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium, magnesium or calcium salts, also ammonium salts derived from ammonia or suitable organic amines. They come into 15 consideration especially aliphatic, cycloaliphatic, cycloaliphatic-aliphatic or araliphatic primary, secondary or tertiary mono-, di- or poly-amines, such as lower alkylamines, for example di- or tri-ethylamine, hydroxy-lower alkylamines, for example 2-hydroxyethylamine, bis(2-hydroxyethyl)amine or tris(2-hydroxyethyl)amine, basic aliphatic esters or carboxylic acids, for example 4-aminobenzoic acid 2-diethylaminoethyl ester, lower alkyleneamines, 20 for example 1-ethylpiperidine, cycloalkylamines, for example dicyclohexylamine, benzylamines, for example N,N'-dibenzylethylenediamine; also heterocyclic bases, for example of the pyridine type, for example pyridine, collidine or quinoline. If several acidic or basic groups are present, mono- or poly-salts can be formed. Compounds according to the invention having an acidic and a basic group may also be in 25 the form of internal salts, i.e. in the form of zwitterions and another part of the molecule in the form of a normal salt.

In the case of the above-mentioned individual compounds the pharmaceutically acceptable salts are included in each case insofar as the individual compound is capable of salt formation.

30 The compounds listed, including the individual compounds mentioned, both in free form and in salt form, may also be in the form of hydrates, or their crystals may include, for example, the solvent used for crystallisation. The present invention relates also to all those forms.

Many of the above-mentioned compounds, including the individual compounds 35 mentioned, contain at least one asymmetric carbon atom. They can therefore occur in the form of R- or S-enantiomers and as enantiomeric mixtures thereof, for example in the form of a racemate. The present invention relates to the use of all those forms and to the use of all further isomers, and of mixtures of at least 2 isomers, for example mixtures of diastereoisomers or enantiomers which can occur when there are one or more further 40 asymmetric centres in the molecule. Also included are, for example, all geometric isomers,

for example cis- and trans-isomers, that can occur when the compounds contain one or more double bonds.

The doses required for the present invention depend entirely on the type of aromatase inhibitor that is used. Some inhibitors are more active than others and hence lower
5 amounts of the former inhibitors could be used. The dosage levels do depend on the patient also.

Typically, the amount of aromatase inhibitor may be selected from amounts that lower estrogen levels to post-menopausal levels in a female. For example the amount of aromatase inhibitor may be selected from amounts that lower the level of estrogen to at or
10 about 100 pmoles/litre or less as measured by standard immunoassay techniques.

The aromatase inhibitor is administered in a single dose in amounts preferably selected from amounts in the range of from at or about 5 mg to at or about 500 mg and the daily doses of follicle stimulating hormone range from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.

15 It may be said that the amount of aromatase inhibitor is selected from amounts that lower estrogen levels to post-menopausal levels in a female. For example, the amount of aromatase inhibitor may be selected from amounts that lower estrogen levels to about 100 pmol/L or less as measured by standard immunoassay techniques. Typically the aromatase inhibitor may be administered in a single dose selected from amounts
20 comprising 10 mg, 20 mg, 25 mg or 30 mg.

Examples of preferred dosages for the single doses are when the aromatase inhibitor is letrozole the single dose administered is from at or about 5 mg to at or about 100 mg. When the AI is anastrozole, it may be administered in a single dose of from at or about 5 mg to at or about 50 mg. When the AI is vorozole, the single dose may be from at or about
25 10 mg to at or about 200 mg. Exemestane is preferably administered in a daily dose of from at or about 50 to at or about 500mg. Preferred is for administration of the single dose to be administered on any of days 1 to 5 of the menstrual cycle. The separate doses AI and FSH may be administered simultaneously, sequentially, separately, or consecutively, with or without a gap.

30 A "POOR RESPONDER" is considered to be a patient who fails to ovulate or who fails to develop an adequate number of pre-ovulatory ovarian follicles after at least one cycle of FSH alone or FSH plus LH at an average daily dose of up to the typical maximum doses, for example, 600 I.U. of F.S.H. for an assisted reproductive technology as described below.

ASSISTED REPRODUCTIVE TECHNOLOGIES (ART)

35 Assisted reproductive technologies (ART) includes, for example, the following techniques:

***In vitro* fertilization (IVF)**, in which oocytes are aspirated from pre-ovulatory follicles, combined with sperm *in vitro* and viable embryos are selected and placed in the uterus.

Gamete Intrafallopian Transfer Procedure (GIFT), in which oocytes and sperm are combined in a catheter, and placed in the fallopian tube, so that conception occurs in the fallopian tube.

Zygote Intrafallopian Transfer Procedure (ZIFT), in which the collected oocytes are
5 combined with sperm, and fertilized embryos are transferred to the fallopian tube.

Intracytoplasmic Sperm Injection (ICSI), in which each oocyte is directly injected with a single sperm via a microscopic needle, and viable embryos are selected for placement in the uterus or the fallopian tube.

Intrauterine Insemination (IUI). Intrauterine insemination (IUI) is a fertility procedure
10 in which motile sperm are washed, concentrated, and injected directly into a woman's uterus.

Therapeutic Donor Insemination (TDI) involves the use of timed insemination of sperm from a donor rather than from the husband.

Controlled ovarian Hyperstimulation (COH) for timed intercourse, for IUI or for other
15 ART procedures such as IVF encompasses the concept of deliberate and regulated induction of superovulation, but also refers to production of a hormonal response intended to lead to the production of multiple eggs in the woman's ovaries and to favor implantation of the embryo into the endometrium.

The female in need of treatment is a preferably a human being, but females of other
20 species could benefit from the treatment of the invention and hence the invention encompasses such applications where applicable.

Pharmaceutical Formulations

The pharmaceutical compositions that can be prepared according to the invention are compositions for enteral, such as peroral or rectal administration, also for transdermal or
25 sublingual administration, and for parenteral, for example intravenous, subcutaneous and intramuscular, administration. Suitable unit dose forms, especially for peroral and/or sublingual administration, for example dragees, tablets or capsules, comprise preferably from approximately 0.01 mg to approximately 20 mg, especially from approximately 0.1 mg to approximately 10 mg, of one of the above-mentioned compounds or of a
30 pharmaceutically acceptable salt thereof, together with pharmaceutically acceptable carriers. The preferred form of administration is oral. The proportion of active ingredient in such pharmaceutical compositions may range from approximately 0.001% to approximately 60%, preferably from approximately 0.1% to approximately 20%.

Suitable excipients for pharmaceutical compositions for oral administration are
35 especially fillers, such as sugars, for example lactose, saccharose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, and binders, such as starches, for example corn, wheat, rice or potato starch, gelatin, tragacanth, methylcellulose and/or hydroxypropylcellulose, disintegrators, such as the above-mentioned starches, also carboxymethyl starch, cross-
40 linked polyvinylpyrrolidone, agar, alginic acid or a salt thereof, such as sodium alginate,

and/or cellulose, for example in the form of crystals, especially in the form of microcrystals, and/or flow regulators and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate, cellulose and/or polyethylene glycol.

5 Dragee cores can be provided with suitable, optionally enteric, coatings, there being used inter alia concentrated sugar solutions which may comprise gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, or coating solutions in suitable solvents or solvent mixtures, or, for the preparation of enteric coatings, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate.

10 Other orally administrable pharmaceutical compositions are dry-filled capsules consisting of gelatin, and also soft sealed capsules consisting of gelatin and a plasticiser, such as glycerol or sorbitol. The dry-filled capsules may contain the active ingredient in the form of granules, for example in admixture with fillers, such as lactose, binders, such as starches, and/or glidants, such as talc or magnesium stearate, and, if desired, stabilisers. In
15 soft capsules, the active ingredient is preferably dissolved or suspended in suitable oily excipients, such as fatty oils, paraffin oil or liquid polyethylene glycols, to which stabilisers and/or anti-bacterial agents may also be added. There may also be used capsules that are easily bitten through, in order to achieve by means of the sublingual ingestion of the active ingredient that takes place as rapid an action as possible.

20 Suitable rectally administrable pharmaceutical compositions are, for example, suppositories that consist of a combination of the active ingredient with a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, paraffin hydrocarbons, polyethylene glycols or higher alkanols. There may also be used gelatin rectal capsules, which contain a combination of the active ingredient with a base material.
25 Suitable base materials are, for example, liquid triglycerides, polyethylene glycols or paraffin hydrocarbons.

Suitable formulations for transdermal administration comprise the active ingredient together with a carrier. Advantageous carriers include absorbable pharmacologically acceptable solvents that serve to facilitate the passage through the skin of the host.

30 Transdermal systems are usually in the form of a bandage that comprises a support, a supply container containing the active ingredient, if necessary together with carriers, optionally a separating device that releases the active ingredient onto the skin of the host at a controlled and established rate over a relatively long period of time, and means for securing the system to the skin.

35 Suitable for parenteral administration are especially aqueous solutions of an active ingredient in water-soluble form, for example in the form of a water-soluble salt, and also suspensions of active ingredient, such as corresponding oily injection suspensions, there being used suitable lipophilic solvents or vehicles, such as fatty oils, for example sesame oil, or synthetic fatty acid esters, for example ethyl oleate, or triglycerides, or aqueous

injection suspensions that comprise viscosity-increasing substances, for example sodium carboxymethylcellulose, sorbitol and/or dextran, and, optionally, stabilisers.

Dyes or pigments may be added to the pharmaceutical compositions, especially to the tablets or dragee coatings, for example for identification purposes or to indicate different
5 doses of active ingredient.

The pharmaceutical compositions of the present invention can be prepared in a manner known per se, for example by means of conventional mixing, granulating, confectioning, dissolving or lyophilising processes. For example, pharmaceutical compositions for oral administration can be obtained by combining the active ingredient
10 with solid carriers, optionally granulating a resulting mixture, and processing the mixture or granules, if desired or necessary after the addition of suitable excipients, to form tablets or dragee cores.

The invention that is claimed is described in detail in the following Example, which is intended merely to illustrate the invention, and in no way to represent a limitation thereof.

15 EXAMPLE

Seven infertile patients (3 with PCOS and 4 with unexplained infertility) undergoing ovarian stimulation and cycle monitoring for intrauterine insemination, received letrozole as a single 20 mg dose on day 3 of the menstrual cycle for 9 treatment cycles. HCG 10,000 IU was given to trigger ovulation. Follicular development was monitored by transvaginal
20 ultrasound and hormonal assays of estrogen and LH. Various parameters of the single dose treatment cycles have been compared with a historical control group which included 105-treatment cycles in which letrozole was administered in a dose of 2.5mg/day, from day 3 to 7 of the menstrual cycle.

Ovulation occurred in 8 out of the 9 single-dose treatment cycles and pregnancy was
25 achieved in one patient. In the historical control group (33 PCOS patients underwent 42 treatment cycles and 51 patients with unexplained infertility underwent 63 cycles), ovulation rate was 83.7% in the PCOS group and pregnancy rate was 13.3 % (14% in the PCOS group and 12.7% in the unexplained infertility group). There was no significant difference in the various characteristics of the single dose treatment cycles from the 5-day treatment
30 cycles of letrozole.

The aromatase inhibitor letrozole has been shown to be effective in stimulating ovarian follicular development. This is believed to be due to decreasing estrogen synthesis without having a direct anti-estrogenic effect, which leads to release of the estrogen negative feedback on the pituitary and/or hypothalamus resulting in an increase in gonadotropin
35 secretion. Administering letrozole as a single dose on day three of the menstrual cycle has the advantage of allowing rapid clearance of the letrozole from the body due to its short half-life (around two days). This leads to negligible levels of letrozole in the body around the ovulation and early embryogenesis period. Besides the increased safety of the single dose administration, it is more convenient.

Table 1		
	Letrozole single-dose cycles	Letrozole 5 days cycle
Day of hCG administration	12.9 (2.9)	13.2 (2.5)
Endometrial thickness on hCG day (cm)	0.9 (0.11)	0.9 (0.2)
Follicles>1.5cm	2.29 (1.3)	1.9 (0.2)
Estradiol on hCG day (pmol/L)	831 (359)	919 (782)
Estradiol/mature follicle (pmol/L)	390 (74)	462 (257)
LH (IU/L)	19.1 (12.7)	18.2 (16.2)

While the invention has been described with particular reference to certain
embodiments thereof, it will be understood that changes and modifications may be made
5 by those of ordinary skill in the art within the scope and spirit of the following claims.

In the claims, the word "comprising" means "including the following elements (in the
body), but not excluding others"; the phrase "consisting of" means "excluding more than
traces of other than the recited ingredients"; and the phrase "consisting essentially of"
means "excluding unspecified ingredients which materially affect the basic characteristics
10 of the composition".

REFERENCES

-
- ¹ Garcia J, Jones GS and Wentz AC. The use of clomiphene citrate. *Fertil Steril* 1977, 28:707-17
- ² Goldfarb A.F., Morales A., Rakoff A.E. and Protos P. Critical review of 160 clomiphene-related pregnancies. *Obstet Gynecol* 1968, 31:342-345.
- ³ Randall JM, Templeton A. Cervical mucus score and in vitro sperm mucus interaction in spontaneous and clomiphene citrate cycles. *Fertil Steril* 1991; 56:465-8; and Gelety TJ and Buyalos RP. The effect of clomiphene citrate and menopausal gonadotropins on cervical mucus in ovulatory cycles. *Fertil Steril* 1993, 60:471-476.
- ⁴ Gonen Y, Casper RF. Sonographic Determination of an Adverse Effect of Clomiphene Citrate on Endometrial Growth. *Hum Reprod* 1990; 5:670-4.
- ⁵ Mikkelsen TJ, Kroboth PD, Cameron WJ, Dittert LW, Chungi V, Manberg PJ. Single-dose pharmacokinetics of clomiphene citrate in normal volunteers. *Fertil Steril* 1986; 46:392-6.
- ⁶ Fluker MR, Urman B, Mackinnon M et al. Exogenous gonadotropin therapy in World Health Organization Groups I and II ovulatory disorders. *Obstet Gynecol* 1994; 83:189-96.
- ⁷ Fisch P, Casper RF, Brown SE, Wrixon W, Collins JA, Reid RL, Simpson C. Unexplained infertility: evaluation of treatment with clomiphene citrate and human chorionic gonadotropin. *Fertil Steril* 1989; 51(5): 828-33.
- ⁸ Sereepapong W, Triratanachai S, Sampatanukul P, Pruksananonda K, Boonkasemsanti K and Reinprayoon D. Effects of clomiphene citrate on the endometrium of regularly cycling women. *Fertil Steril* 2000; (73): 287-91.
- ⁹ Taymor M L. The regulation of follicle growth: some clinical implications in reproductive endocrinology. *Fertil Steril* 1996; 65(2): 235-47
- ¹⁰ Wysowski DE. Use of fertility drugs in the United States, 1979 through 1991. *Fertil Steril* 1993;60:1096-98
- ¹¹ Adashi: Clomiphene citrate: mechanism(s) and site(s) of action—a hypothesis revisited. *Fertil Steril* 1984; 42(3):331-44
- ¹² Dickey RP, Vorys N, Stevens VC, Besch PK, Hamwi GJ, Ullery JC. Observations on the mechanism of action of clomiphene (MRL-41). *Fertil Steril* 1965;16:485-94

-
- ¹³ Kettel LM, Roseff SJ, Berga SL, Mortola JF, Yen SS. Hypothalamic-pituitary-ovarian response to clomiphene citrate in women with polycystic ovary syndrome. *Fertil Steril* 1993;59(3):532-8.
- ¹⁴ Archer DF, Hofmann G, Brzyski R, Ross B, Scott RT, Philput CB, Oehninger S, Ware JC. Effects of clomiphene citrate on episodic luteinizing hormone secretion throughout the menstrual cycle. *Am J Obstet Gynecol* 1989;161(3):581-9.
- ¹⁵ Adashi: Clomiphene citrate: mechanism(s) and site(s) of action--a hypothesis revisited. *Fertil Steril* 1984; 42(3):331-44
- ¹⁶ Adashi: Clomiphene citrate: mechanism(s) and site(s) of action--a hypothesis revisited. *Fertil Steril* 1984; 42(3):331-44
- ¹⁷ Yagel S, Ben-Chetrit A, Anteby E, Zacut D, Hochner-Celnikier D and Ron M. The effect of ethinyl estradiol on endometrial thickness and uterine volume during ovulation induction by clomiphene citrate. *Fertil Steril* 1992, 57:33-36.
- ¹⁸ Ben-Ami M, Geslevich Y, Matilsky M, Battino S, Weiner E et al. Exogenous estrogen therapy concurrent with clomiphene citrate--lack of effect on serum sex hormone levels and endometrial thickness. *Gynecol Obstet Invest.* 1994;37(3):180-2.
- ¹⁹ Bateman BG, Nunley WC Jr and Kolp LA. Exogenous estrogen therapy for treatment of clomiphene citrate--induced cervical mucus abnormalities: is it effective? *Fertil Steril* 1990, 54:577-9
- ²⁰ Wu CH, Winkel CA. The effect of therapy initiation day on clomiphene citrate therapy. *Fertil Steril* 1989;52:564-568
- ²¹ Saleh A, Biljan MM, Tan SSSL and Tulandi T. Effects of Tamoxifen (Tx) on Endometrial Thickness and Pregnancy Rates in Women Undergoing Superovulation with Clomiphene Citrate (CC) and Intrauterine Insemination (IUI). *Fertil Steril* 2000; 74(S1):S90
- ²² Sioufi A, Gauducheau N, Pineau V, et al: Absolute bioavailability of letrozole in healthy post-menopausal women. *Biopharm Drug Dispos* 1997; 18:779-89; and Sioufi A, Sandrenan N, Godbillon J, Trunet P, Czendlik C, Howald H, Pfister C, Ezzet F. Comparative bioavailability of letrozole under fed and fasting conditions in 12 healthy subjects after a 2.5 mg single oral administration. *Biopharm Drug Dispos* 1997; 18(6): 489-97.
- ²³ Cole PA, Robinson CH. Mechanism and inhibition of cytochrome P-450 aromatase, *J. Med. Chem.* 33 (1990) 2933-2944; and Akhtar M, Njar VCO, Wright JN. Mechanistic

studies on aromatase and related C-C bond cleaving P-450 enzymes, *J Steroid Biochem Mol Biol* 1993; (44): 375-387.

²⁴ Santen RJ, Manni A, Harvey H, Redmond C. Endocrine treatment of breast cancer in women. *Endocrine Rev* 1990;11:1-45.

²⁵ Nebert DW, Nelson DR, Coon MJ, Estabrook RW, Feyereisen R, et al. The P-450 superfamily: update on new sequences, gene mapping and recommended nomenclature, *DNA Mol. Biol.* 10 (1991) 1-14.

²⁶ R.C. Coombes, P. Goss, M. Dowsett, J.C. Gazet, A.M.H. Brodie, 4-Hydroxyandrostenedione treatment of postmenopausal patients with advanced breast cancer, *The Lancet* 2,1984,1237-1239.

²⁷ Santen RJ, Lipton A, Kendall J. Successful medical adrenalectomy with aminoglutethimide: role of altered drug metabolism. *J Am Med Assoc* 1974;230:1661.

²⁸ Harris AL, Dowsett M, Smith IE et al. Aminoglutethimide in premenopausal patients with breast cancer: endocrine studies and tumour response. *Cancer Chemother Pharmacol* 1980;5:23.

²⁹ Marty M, Gershanovich M, Campos B, Romien G, Lurie H, Bonaventura T, et al. Als, a new potent, selective aromatase inhibitor superior to aminoglutethimide (AG) in postmenopausal women with advanced breast cancer previously treated with antiestrogens. *Proc Am Soc Clin Oncol* 1997;16:156; and Dowsett M. Biological background to aromatase inhibition. *The Breast* 1996; 5:196-201.

³⁰ Sinha S, Kaseta J, Santner LM et al: Effects of CGS20267 on ovarian aromatase and gonadotropin levels in the rat. *Breast Cancer Res Treat* 1998; 48:45-51.

³¹ Shetty G, Krishnamurthy H, Krishnamurthy HN et al: Effect of estrogen deprivation on the reproductive physiology of male and female primates. *J Steroid Biochem Biol* 1997; 61:157-66.

³² Geisler J, King N, Dowsett M et al. Influence of anastrozole (Arimidex), a selective, non-steroidal aromatase inhibitor, on in vivo aromatisation and plasma estrogen levels in postmenopausal women with breast cancer. *Br J Cancer* 1996; 74: 1286-91.

³³ Sioufi A, Gauducheau N, Pineau V, et al: Absolute bioavailability of letrozole in healthy post-menopausal women. *Biopharm Drug Dispos* 1997; 18:779-89.

-
- ³⁴ Trunet PF, Mueller P, Bhatnagar PS, Dickes I, Monnet G and White G. Open dose-finding study of a new potent and selective nonsteroidal aromatase inhibitor, CGS 20 267, in healthy male subjects. *J Clin Endocrinol Metab* 1993; 77: 319-323.
- ³⁵ *Femara professional information brochure*. Novartis Pharmaceuticals Corporation. East Hanover, New Jersey 2000.
- ³⁶ Mitwally MFM and Casper RF. The Use of an Aromatase Inhibitor for Induction of Ovulation in Cases of Clomiphene Citrate Failure. [Abstract number O-178] In: Program and abstracts of The 16th Annual Meeting of the European Society for Human Reproduction and Embryology (ESHRE), June 2000, Bologna, Italy.
- (2) Mitwally MFM and Casper RF. Aromatase Inhibition: A novel Method of Ovulation Induction in Women With Polycystic Ovarian Syndrome *Reprod Technol* 2000; 10(5) In Press.
- ³⁷ Mitwally MFM and Casper RF. The Aromatase Inhibitor, Letrozole: a Promising Alternative for Clomiphene Citrate for Induction of Ovulation. [Abstract number O-091] In: Program and abstracts of The 56th Annual Meeting of the American Society for Reproductive Medicine (ASRM), October 2000, San Diego, CA, USA. Winner of the Prize paper award of the Society of Reproductive Endocrinology and Infertility; and Mitwally MFM and Casper RF: Use of an aromatase inhibitor for induction of ovulation in patients with an inadequate response to clomiphene citrate. *Fertil Steril* 2001; 75(2): 305-9.
- ³⁸ Mitwally MFM and Casper RF. The Aromatase Inhibitor, Letrozole, Decreases FSH Dose Required for Ovarian Superovulation. The 46th Annual Meeting of the Canadian Fertility and Andrology Society. Newfoundland, Canada. September 2000; and Mitwally MFM and Casper RF. Aromatase inhibition decreases FSH dose needed during controlled ovarian hyperstimulation: A controlled prospective trial. Meeting of the Society for Gynecologic Investigation, March 2001, Toronto, Canada. Winner of the President Presenter's Award
- ³⁹ Mitwally MFM and Casper RF. Aromatase Inhibition Improves Ovarian Response to FSH: A Potential Option for Low Responders During Ovarian Stimulation. The 48th meeting of the Pacific Coast Fertility Society Meeting, Rancho Las Palmas Resort and Spa, CA, USA. April 2001.
- ⁴⁰ Weil S; Vendola K; Zhou J; Bondy CA. Androgen and follicle-stimulating hormone interactions in primate ovarian follicle development. *J Clin Endocrinol Metab* 1999;84(8):2951-6.
- ⁴¹ Vendola KA, Zhou J, Adesanya OO, Weil SJ, Bondy CA. Androgens stimulate early stages of follicular growth in the primate ovary. *J Clin Invest* 1998;101(12):2622-9.

CLAIMS

1. Use of an aromatase inhibitor for the preparation of a medicament for inducing ovulation in a female suffering from anovulatory infertility, wherein the medicament is administered as a single dose of at least one aromatase inhibitor.
2. Use as claimed in Claim 1, wherein the aromatase inhibitor is selected from aromatase inhibitors having a half-life of from at or about 8 hours to at or about 4 days.
3. Use as claimed in Claim 1, wherein the aromatase inhibitor is selected from aromatase inhibitors having a half-life of about 2 days.
4. Use as claimed in any one preceding claim, wherein the aromatase inhibitor is selected from non-steroidal and reversible aromatase inhibitors.
5. Use as claimed in any one preceding claim, wherein the aromatase inhibitor is administered orally.
6. Use as claimed in any one preceding claim, wherein the amount of aromatase inhibitor is selected from amounts that lower estrogen levels to post-menopausal levels in a female.
7. Use as claimed in any one preceding claim, wherein the amount of aromatase inhibitor is selected from amounts that lower estrogen levels to at or about 100 pmol/L or less as measured by standard immunoassay techniques.
8. Use as claimed in any one preceding claims, wherein the aromatase inhibitor is administered in a single dose selected from amounts in the range of from at or about 5 mg to at or about 500 mg.
9. Use as claimed in any one preceding claim, wherein the aromatase inhibitor is administered in a single dose selected from amounts comprising 5 mg, 10 mg, 20 mg, 25 mg or 30 mg.
10. Use as claimed in any one of claims 1 to 8, wherein the aromatase inhibitor is letrozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg.
11. Use as claimed in any one of claims 1 to 8, wherein the aromatase inhibitor is anastrozole and is administered in a single dose of from at or about 5 mg to at or about 50 mg.
12. Use as claimed in any one of claims 1 to 8, wherein the aromatase inhibitor is anastrozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg.
13. Use as claimed in any one of claims 1 to 8, wherein the aromatase inhibitor is vorozole and is administered in a single dose of from at or about 10 mg to at or about 200 mg.
14. Use as claimed in any one of claims 1 to 8, wherein the aromatase inhibitor is vorozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg.
15. Use as claimed in any one of claims 1 to 8, wherein the aromatase inhibitor is exemestane and is administered in a single dose of from at or about 50 mg to at or about 500 mg.

16. Use as claimed in any one of claims 1 to 8, wherein the aromatase inhibitor is exemestane and is administered in a single dose of from at or about 25 mg to at or about 500 mg
17. Use as claimed in any one preceding claim, wherein the medicament is administered on any one of days 1 to 5 of the menstrual cycle.
18. Use according to any one preceding claim, wherein the medicament is administered on day 3 of the menstrual cycle.
19. Use of an aromatase inhibitor for the preparation of a medicament for augmenting ovulation in an ovulating female suffering from unexplained infertility or other types of ovulatory infertility, wherein the medicament is administered to said female as a single dose of at least one aromatase inhibitor (AI) early in one or more menstrual cycles.
20. Use as claimed in Claim 19, wherein the medicament is administered on any of days 1 to 5 of the menstrual cycle.
21. Use as claimed in Claim 19, wherein the medicament is administered on day 3 of the menstrual cycle.
22. Use as claimed in Claim 19, 20 or 21, wherein the aromatase inhibitor is selected from aromatase inhibitors having a half-life of about 8 hours to about 4 days.
23. Use as claimed in Claim 19, 20 or 21, wherein the aromatase inhibitor is selected from aromatase inhibitors having a half-life of about 2 days.
24. Use as claimed in any one of claims 19 to 23, wherein the aromatase inhibitor is selected from non-steroidal and reversible aromatase inhibitors.
25. Use as claimed in any one of claims 19 to 24, wherein the aromatase inhibitor is administered orally.
26. Use as claimed in any one of claims 19 to 25, wherein the amount of aromatase inhibitor is selected from amounts that lower estrogen levels to post-menopausal levels in a female.
27. Use as claimed in any one of claims 19 to 25, wherein the amount of aromatase inhibitor is selected from amounts that lower estrogen levels to at or about 100 pmoles/litre or less as measured by standard immunoassay techniques.
28. Use as claimed in any one of claims 19 to 27, wherein the aromatase inhibitor is administered in a single dose selected from amounts in the range of from at or about 5 mg to at or about 500 mg.
29. Use as claimed in any one of claims 19 to 28, wherein the aromatase inhibitor is administered in a single dose selected from amounts comprising 5 mg, 10 mg, 20 mg, 25 mg or 30 mg.
30. Use as claimed in any one of claims 19 to 28, wherein the aromatase inhibitor is letrozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg.
31. Use as claimed in any one of claims 19 to 28, wherein the aromatase inhibitor is anastrozole and is administered in a single dose of from about 5 mg to about 50 mg.

32. Use as claimed in any one of claims 19 to 28, wherein the aromatase inhibitor is anastrozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg
33. Use as claimed any one of claims 19 to 28, wherein the aromatase inhibitor is vorozole and is administered in a single dose of from about 10 mg to about 200 mg.
34. Use as claimed in any one of claims 19 to 28, wherein the aromatase inhibitor is vorozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg.
35. Use as claimed in any one of claims 19 to 28, wherein the aromatase inhibitor is exemestane and is administered in a single dose of from at or about 25 mg to at or about 500 mg.
36. Use as claimed in any one of claims 19 to 28, wherein the aromatase inhibitor is exemestane and is administered in a single dose of from at or about 50 mg to at or about 500 mg.
37. Use as claimed in any one of claims 19 to 36, wherein the medicament is administered on any one of days 1 to 5 of the menstrual cycle.
38. Use according to any one of claims 19 to 36, wherein the medicament is administered on day 3 of the menstrual cycle.
39. Use of an aromatase inhibitor for the preparation of a medicament for use with follicle stimulating hormone (FSH) in the treatment of an anovulatory female for infertility, wherein the medicament is administered as single dose.
40. Use as claimed in Claim 39, wherein separate doses AI and FSH are administered simultaneously, separately or sequentially.
41. Use as claimed in Claim 39 or 40, wherein the aromatase inhibitor is selected from aromatase inhibitors having a half-life of about 8 hours to about 4 days.
42. Use as claimed in Claim 39, 40 or 41, wherein the aromatase inhibitor is selected from aromatase inhibitors having a half-life of about 2 days.
43. Use as claimed in any one of Claims 39 to 42, wherein the aromatase inhibitor is selected from non-steroidal and reversible aromatase inhibitors.
44. Use as claimed in any one of Claims 39 to 43, wherein the aromatase inhibitor is administered orally.
45. Use as claimed in any one of Claims 38 to 44, wherein the amount of aromatase inhibitor is selected from amounts that lower estrogen levels to post-menopausal levels in a female.
46. Use as claimed in any one of Claims 38 to 45, wherein the amount of aromatase inhibitor is selected from amounts that lower estrogen levels to at or about 100 pmoles/litre or less as measured by standard immunoassay techniques.
47. Use as claimed in any one of Claims 38 to 46, wherein the aromatase inhibitor is administered in a dose selected from amounts in the range of from at or about 5 mg to at or about 500 mg and wherein the daily doses of follicle stimulating hormone ranges

- from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
48. Use as claimed in any one of Claims 38 to 47, wherein the aromatase inhibitor is administered in a dose selected from amounts comprising 5 mg, 10 mg, 20 mg, 25 mg or 30 mg.
 49. Use as claimed in any one of Claims 38 to 46, wherein the aromatase inhibitor is letrozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
 50. Use as claimed in any one of Claims 38 to 46, wherein the aromatase inhibitor is anastrozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
 51. Use as claimed in any one of Claims 38 to 46, wherein the aromatase inhibitor is vorozole and is administered in a single dose of from at or about 10 mg to at or about 200 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
 52. Use as claimed in any one of Claims 38 to 46, wherein the aromatase inhibitor is exemestane and is administered in a single dose of from at or about 50 mg to at or about 500 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
 53. Use as claimed in any one of Claims 38 to 46, wherein the aromatase inhibitor is vorozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
 54. Use as claimed in any one of Claims 38 to 46, wherein the aromatase inhibitor is exemestane and is administered in a single dose of from at or about 5 mg to at or about 100 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
 55. Use as claimed in any one of claims 38 to 54, wherein the medicament is administered on any one of days 1 to 5 of the menstrual cycle.
 56. Use according to claim 55, wherein the medicament is administered on day 3 of the menstrual cycle.

57. Use of a single dose of an aromatase inhibitor (AI) for the preparation of a medicament for use with follicle stimulating hormone (FSH) in the treatment of a female for infertility, wherein the female is considered to be a poor responder to FSH, wherein the medication is administered as a single dose.
58. Use as claimed in claim 57, wherein the doses of AI and FSH are administered simultaneously, separately or sequentially.
59. Use as claimed in Claim 57 or 58, wherein the aromatase inhibitor is selected from aromatase inhibitors having a half-life of about 8 hours to about 4 days.
60. Use as claimed in Claim 57 or 58, wherein the aromatase inhibitor is selected from aromatase inhibitors having a half-life of about 2 days.
61. Use as claimed in any one of Claims 57 to 60 wherein the aromatase inhibitor is selected from non-steroidal and reversible aromatase inhibitors.
62. Use as claimed in any one of Claims 57 to 61 wherein the aromatase inhibitor is administered orally.
63. Use as claimed in any one of Claims 57 to 62 wherein the amount of aromatase inhibitor is selected from amounts that lower estrogen levels to post-menopausal levels in a female.
64. Use as claimed in any one of Claims 57 to 62, wherein the amount of aromatase inhibitor is selected from amounts that lower estrogen levels to about 100 pmoles/litre or less as measured by standard immunoassay techniques.
65. Use as claimed in any one of Claims 57 to 63, wherein the aromatase inhibitor is administered in a dose selected from amounts in the range of from at or about 5 mg to at or about 500 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
66. Use as claimed in any one of Claims 57 to 65, wherein the aromatase inhibitor is administered in a dose selected from amounts comprising 5 mg, 10 mg, 20 mg, 25 mg or 30 mg.
67. Use as claimed in any one of Claims 57 to 65, wherein the aromatase inhibitor is letrozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
68. Use as claimed in any one of Claims 57 to 65, wherein the aromatase inhibitor is anastrozole and is administered in a single dose of from at or about 5 mg to at or about 50 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
69. Use as claimed in any one of Claims 57 to 65, wherein the aromatase inhibitor is vorozole and is administered in a single dose of from at or about 10 mg to at or about

200 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.

70. Use as claimed in any one of Claims 57 to 65, wherein the aromatase inhibitor is exemestane and is administered in a single dose of from at or about 50 mg to at or about 500 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
71. Use as claimed in any one of Claims 57 to 65, wherein the aromatase inhibitor is anastrozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
72. Use as claimed in any one of Claims 57 to 65, wherein the aromatase inhibitor is vorozole and is administered in a single dose of from at or about 5 mg to at or about 100 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
73. Use as claimed in any one of Claims 57 to 65, wherein the aromatase inhibitor is exemestane and is administered in a single dose of from at or about 50 mg to at or about 500 mg and wherein the daily doses of follicle stimulating hormone ranges from at or about 25 to at or about 600 units or an equivalent dosage in another form of administration.
74. Use as claimed in any one preceding claim, wherein the aromatase inhibitor is administered on any one of days 1 to 5 of the menstrual cycle.
75. Use as claimed in any one preceding claim, wherein the aromatase inhibitor is administered on day 3 of the menstrual cycle.
76. A kit for treating infertility in a female comprising a single dose of an aromatase inhibitor together with a pharmaceutically acceptable carrier in combination with a plurality of daily doses of follicle stimulating hormone together with a pharmaceutically acceptable carrier.
77. A pharmaceutical package, comprising a single dose of an aromatase inhibitor, and instructions for its use in ovulation induction.
78. A pharmaceutical package, comprising a single dose of an aromatase inhibitor, a plurality of doses of FSH, and instructions for its use in ovulation induction in a female patient.
79. A pharmaceutical package according to claim 78, wherein the female patient is a poor responder to FSH.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 02/00527

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61P15/00 A61P5/24 A61K31/56 A61K31/4196

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS, PHARMAPROJECTS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	MITWALLY MOHAMED F ET AL: "Aromatase inhibition for ovarian stimulation: future avenues for infertility management." CURRENT OPINION IN OBSTETRICS & GYNECOLOGY. ENGLAND JUN 2002, vol. 14, no. 3, June 2002 (2002-06), pages 255-263, XP001084791	1-14, 17-34, 37-51, 53, 55-69, 71,72, 74,75
P,Y	ISSN: 1040-872X the whole document	15,16, 35,36, 52,54, 70,73, 76-79
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

13 August 2002

Date of mailing of the international search report

12/09/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Giacobbe, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 02/00527

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	MITWALLY MOHAMED FAROUK M ET AL: "Aromatase inhibition improves ovarian response to follicle-stimulating hormone in poor responders(1)." FERTILITY AND STERILITY. UNITED STATES APR 2002, vol. 77, no. 4, April 2002 (2002-04), pages 776-780, XP001088750 ISSN: 0015-0282 the whole document	1-79
Y	"The ovulatory cycle and drug therapy - Letrozole Protocol" INTERNET, 'Online! February 2001 (2001-02), XP002205541 Retrieved from the Internet: <URL:http://www.baby-makers.com/theovulatorycycleanddrugtherapy.html> 'retrieved on 2002-07-10! the whole document	1-79
Y	MITWALLY M.F.M. AND CASPER R.F.: "Use of aromatase inhibitor for induction of ovulation in patients with an inadequate response to clomiphene citrate" FERTILITY AND STERILITY, vol. 75, no. 2, February 2001 (2001-02), pages 305-309, XP001088751 us the whole document page 308, column 2, line 2 - line 14	1-79
Y	MITWALLY M F ET AL: "The use of an aromatase inhibitor for induction of ovulation in cases of clomiphene citrate failure." HUMAN REPRODUCTION (OXFORD), vol. 15, no. Abstract Book 1, June 2000 (2000-06), pages 71-72, XP001094320 16th Annual Meeting of the European Society of Human Reproduction and Embryology; Bologna, Italy; June 25-28, 2000 ISSN: 0268-1161 the whole document	1-79

-/--

INTERNATIONAL SEARCH REPORT

In ational Application No

PCT/CA 02/00527

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with Indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Mitwally MFM et al. 'The Aromatase Inhibitor Letrozole: a Promising Alternative for Clomiphene Citrate for Induction of Ovulation' Abstract No. 0-91 in: Program and Abstracts of the 56th Annual Meeting of the American Society for Reproductive Medicine (ASRM), October 2000, San Diego (CA), USA XP001088806 cited in the application the whole document ---	1-79
Y	MITWALLY, MFM, ET AL.: "Aromatase Inhibition Decreases FSH dose Needed During Controlled Ovarian Hyperstimulation: a Controlled Perspective Trial" J. SOC. GYNECOL. INVESTIG., vol. 8, no. 1(Supplement), February 2001 (2001-02), page 85A XP001084377 cited in the application the whole document ---	1-79
Y	MITWALLY, MFM., ET AL.: "Aromatase inhibition: a novel method of ovulation induction in women with polycystic ovary syndrome" REPRODUCTIVE TECHNOLOGIES, vol. 10, no. 5, 2000, pages 244-247, XP001084440 the whole document ---	1-79
Y	Mitwally, MFM, et al., 'Aromatase Inhibition Improves Ovarian Response to FSH: a Potential Option for Low Responders During Ovarian Stimulation', Abstract No. o-9 in Abstracts of the 48th Meeting of the Pacific Coast Fertility Society, April 2001, Rancho Las Palmas Resort and Spa, CA, USA XP002207747 cited in the application the whole document ---	1-79
Y	COOMBES R C ET AL: "AROMATASE INHIBITORS AND THEIR USE IN THE SEQUENTIAL SETTING" ENDOCRINE-RELATED CANCER, JOURNAL OF ENDOCRINOLOGY LTD., BRISTOL, GB, vol. 6, no. 2, June 1999 (1999-06), pages 259-263, XP000986876 ISSN: 1351-0088 abstract --- -/--	15,16, 35,36, 52,54, 70,73, 76-79

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 02/00527

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SIOUFI A ET AL: "Absolute bioavailability of letrozole in healthy postmenopausal women." BIOPHARMACEUTICS & DRUG DISPOSITION. ENGLAND DEC 1997, vol. 18, no. 9, December 1997 (1997-12), pages 779-789, XP002209706 ISSN: 0142-2782 abstract page 780, paragraph 2 figure 2 table 1 page 787, line 5 -page 788, line 14	1-79
X	----- DATABASE PHARMAPROJECTS 'Online! PJB Publications Ltd., UK; "Exemestane" Database accession no. 4447 XP002209732 the whole document	77
X	----- DATABASE PHARMAPROJECTS 'Online! PJB Publications Ltd., UK; "Letrozole" Database accession no. 2585 XP002209733 the whole document -----	77